

CONDITIONS FOR ACTIVATION OF GROUP I METABOTROPIC GLUTAMATE RECEPTORS IN RAT HIPPOCAMPUS

DISSERTATION

ZUR

ERLANGUNG DER NATURWISSENSCHAFTLICHEN DOKTORWÜRDE
(DR. SC. NAT.)

VORGELEGT DER

MATHEMATISCH-NATURWISSENSCHAFTLICHEN FAKULTÄT

DER

UNIVERSITÄT ZÜRICH

VON

WEI FAN

AUS

CHINA

PROMOTIONSKOMITEE

PROF. DR. STEPHAN NEUHAUSS
PROF. DR. URS GERBER
PROF. DR. PETER SONDEREGGER

ZÜRICH 2010

The unity of knowledge and action

Wang Yangming

Dedicated to my parents

Zusammenfassung

Metabotrope Glutamatrezeptoren (mGluRs) gehören zur Familie der G-Protein-gekoppelten Rezeptoren und spielen eine wichtige Rolle im Rahmen synaptischer Plastizität, Entwicklung des Nervensystems und Neurodegeneration. mGluRs der Klasse 1 (mGluR1 und mGluR5) können selektiv durch 3,5-Dihydroxyphenylglycin (3,5-DHPG) aktiviert werden. Klasse 1 mGluRs können im Gehirn sowohl prä- als auch postsynaptisch lokalisiert sein; am häufigsten kommen sie in der postsynaptischen Dichte asymmetrischer, glutamaterger Synapsen vor.

Ziel der vorliegenden Arbeit ist herauszufinden, unter welchen Bedingungen Klasse 1 mGluRs im Hippocampus der Ratte aktiviert werden.

In Kapitel 2 zeige ich, dass Klasse 1 mGluRs in CA1 Pyramidenzellen hippocampaler Schnittkulturen nicht tonisch durch basale Glutamatlevel aktiviert werden.

In Kapitel 3 lege ich dar, dass die Aktivierung einer einzelnen CA3 Pyramidenzelle kein mGluR-LTP im direkt mit ihr verbundenen CA1 Pyramidalneuron induziert. Anhand von Paarableitungen mit extrazellulärer Stimulation lässt sich schliessen, dass schätzungsweise 8 aktive CA3-Eingänge auf eine CA1 Pyramidenzelle konvergieren müssen um in dieser mGluR-LTP auszulösen.

In Kapitel 4 zeige ich, dass die Aktivierung einer einzelnen CA3 Pyramidenzelle keinen durch Klasse 1 mGluRs vermittelten somatischen Strom in einem synaptisch verbundenen Interneuron hervorruft. Klasse 1 mGluRs in Interneuronen des Hippocampus werden durch Theta-Oszillationen aktiviert und modulieren deren Frequenz.

Summary

Metabotropic glutamate receptors (mGluRs) are a family of G-protein-coupled receptors and have been shown to play an important role in synaptic plasticity, neuronal development, and neurodegeneration. Group I mGluRs (mGluR1 and mGluR5) are selectively activated by 3,5-dihydroxyphenylglycine (3,5-DHPG). Although Group I mGluRs can be located at both presynaptic and postsynaptic sites in the brain, they are mostly observed at the periphery of the postsynaptic densities of asymmetrical glutamatergic synapses

The aim of my thesis project is to determine the conditions required for activation of group I mGluRs in rat hippocampus.

In chapter 2, I show that group I mGluRs in CA1 pyramidal cells are not tonically activated by ambient glutamate in hippocampal slice cultures.

In chapter 3, I show that activation of a single CA3 pyramidal cell does not induce mGluR-LTD in a connected CA1 pyramidal cell. An estimate based on data obtained from paired recording and extracellular stimulation indicates that 8 active CA3 inputs are required to induce mGluR-LTD in a targeted CA1 pyramidal cell.

In chapter 4, I show that activation of a single CA3 pyramidal cell does not evoke a somatic current mediated by group I mGluRs in a synaptically connected interneuron. I examine the roles of group I mGluRs in hippocampal interneurons during network oscillations and find that their activation modulates theta frequency.

Table of contents

Zusammenfassung.....	4
Summary.....	5
Table of contents.....	6
Chapter 1	
Introduction.....	9
1.1 Ionotropic glutamate receptors.....	10
1.1.1 NMDA receptors.....	11
1.1.2 AMPA receptors.....	12
1.1.3 Kainate receptors.....	13
1.2 Metabotropic glutamate receptors.....	14
1.2.1 Classification.....	14
1.2.2 Structure.....	16
1.2.3 Function.....	17
1.3 Group I metabotropic glutamate receptors.....	19
1.3.1 Distribution of group I mGluRs in the central nervous system.....	19
1.3.2 Subcellular localization of group I mGluRs.....	20
1.3.3 Function of group I mGluRs.....	21
1.4 Long-term potentiation and long-term depression.....	23
1.4.1 Long-term potentiation.....	23
1.4.2 Long-term depression.....	24
1.5 Oscillations.....	26
1.5.1 Function of oscillations.....	26
1.5.2 Oscillations in hippocampus.....	26
1.5.3 Methods of inducing oscillation <i>in vitro</i>	26
1.5.4 Mechanisms of the generation of theta rhythm.....	27
Chapter 2	
Group I metabotropic glutamate receptors in CA1 pyramidal cells are not tonically activated by ambient glutamate in hippocampal slice cultures.....	29
2.1 Abstract.....	30

2.2 Introduction.....	30
2.3 Materials and Methods.....	31
2.3.1 Slice cultures and electrophysiology.....	31
2.3.2 Induction of NMDA currents.....	32
2.3.3 Drugs.....	32
2.4 Results.....	33
2.4.1 Group I mGluRs in CA1 pyramidal cells are not tonically activated by ambient glutamate in hippocampal slice cultures.....	33
2.5 Discussion.....	35
Chapter 3	
Activation conditions for the induction of metabotropic glutamate receptor- dependent LTD in hippocampal CA1 pyramidal cells.....	36
3.1 Abstract.....	37
3.2 Introduction.....	37
3.3 Materials and Methods.....	38
3.3.1 Slice cultures and electrophysiology.....	38
3.3.2 Extracellular stimulation.....	39
3.3.3 Perforated patch-clamp.....	40
3.3.4 Drugs.....	40
3.4 Results.....	41
3.4.1 A single CA3 pyramidal cell does not induce mGluR-LTD in a connected CA1 pyramidal cell.....	41
3.4.2 A single CA3 pyramidal cell does not evoke a somatic current mediated by mGluRs in a connected CA1 pyramidal cell.....	46
3.4.3 Estimating the number of Schaffer collaterals required to induce mGluR-mediated responses.....	49
3.5 Discussion.....	53
Chapter 4	
Group I metabotropic glutamate receptors modulate hippocampal rhythmic activity.....	56
4.1 Abstract.....	57

4.2 Introduction.....	57
4.3 Materials and Methods.....	58
4.3.1 Slice cultures and electrophysiology.....	58
4.3.2 Extracellular stimulation.....	59
4.3.3 Paired recordings.....	59
4.3.4 Drugs.....	59
4.4 Results.....	60
4.4.1 Activation of a single CA3 pyramidal cell does not evoke a somatic current mediated by group I mGluRs in a synaptically connected interneuron.....	60
4.4.2 Extracellular stimulation of Schaffer collaterals evokes group I mGluR-mediated EPSCs in interneurons of the CA1 region.....	62
4.4.3 Group I mGluRs modulate hippocampal theta rhythm.....	63
4.5 Discussion.....	65
Chapter 5	
Conclusion and perspectives.....	66
5.1 Conclusion and discussion.....	67
5.2 Perspectives.....	69
References.....	71
List of abbreviations.....	107
Curriculum Vitae.....	109
Acknowledgements.....	110

Chapter 1

Introduction

L-Glutamate is the major excitatory neurotransmitter in the mammalian central nervous system (CNS) and it acts through a variety of ionotropic (ligand-gated cation channels) and metabotropic (G-protein coupled) receptors (Fig. 1.1, adapted from Kew and Kemp, 2005).




Figure 1.1 Glutamate receptors. Adapted from Kew and Kemp, 2005.

1.1 Ionotropic glutamate receptors

Ionotropic glutamate receptors (iGluRs) are ion channels permeable to cations and are subdivided into *N*-methyl-D-aspartate (NMDA), α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA) and kainate receptors based on agonist preference (Bettler and Mulle, 1995; Dingledine 1999; Kew and Kemp, 2005; Mayer, 2005; Köhr, 2006; Pinheiro and Mulle, 2006; Sprengel, 2006).

iGluRs are tetramers with each subunit possessing an extracellular amino terminal domain, followed by a first transmembrane domain and then a pore forming membrane-residing domain that does not cross the membrane but forms a re-entrant loop entering from and exiting to the cytoplasm. The second and third transmembrane domains are linked by a large extracellular loop and the




Figure 1.2 Schematic drawing of an Ionotropic glutamate receptor subunit. Taken from Kew and Kemp, 2005.

third transmembrane domain is followed by an intracellular carboxy-terminus. The structure of the iGluR ligand binding domain is located in a pocket formed between the extracellular amino terminus (S1 domain) and the extracellular loop between transmembrane domains 3 and 4 (S2 domain) (Fig. 1.2) (Dingledine et al. 1999; Mayer and Armstrong 2004; Kew and Kemp, 2005).

1.1.1 NMDA receptors

NMDA receptors (NMDARs) play pivotal roles in synaptic plasticity, synapse formation and neuronal cell death. A unique property of the NMDAR is its voltage-dependent activation, a result of ion channel block by extracellular Mg^{2+} ions at physiological concentrations. NMDARs require the co-agonist glycine or D-serine. Activation of the NMDAR results in the opening of an ion channel that is nonselective to cations, which allows voltage-dependent flow of Na^{+} and small

amounts of Ca^{2+} ions into the cell and K^{+} out of the cell. Seven NMDAR subunits have been identified: NR1, NR2A, NR2B, NR2C, NR2D, NR3A and NR3B. There are eight different NR1 subunits generated by alternative splicing from a single gene. All NMDARs are heteromeric assemblies composed of two NR1 subunits in combination with at least one type of NR2. The NR3 subunit does not form functional receptors alone, but can co-assemble with NR1/NR2 complexes. NMDAR functional properties are determined by subunit composition (Cotman and Monaghan, 1988; Daw et al., 1993; Mori and Mishina, 1995; Dingledine 1999; Cull-Candy et al., 2001; Debanne et al., 2003; Riedel et al., 2003; Wenthold et al., 2003; Mayer, 2005; Groc and Choquet, 2006; Köhr, 2006; Lau and Zukin, 2007; Paoletti and Neyton, 2007).

1.1.2 AMPA receptors

AMPA receptors (AMPA receptors) are the principal channels that mediate fast excitatory synaptic transmission. AMPARs are composed of four types of subunits, which combine to form tetramers. These subunits are designated as GluR1, GluR2, GluR3 and GluR4, alternatively called GluRA, GluRB, GluRC and GluRD. Most AMPA receptors contain the GluR2 subunit. The AMPAR is permeable to calcium and other cations, such as sodium and potassium. The AMPAR's permeability to calcium is governed by the GluR2 subunit. Functional AMPA receptors are located predominantly in the postsynapse but are also found at extrasynaptic sites and occasionally in the presynapse. The glutamate-mediated transmission efficiency of synaptic AMPARs is influenced by their subunit composition, post-transcriptional and post-translational modifications, the number of synaptic AMPARs, and auxiliary proteins. AMPAR function is regulated over long periods (days to weeks) in response to chronic changes in the level of neuronal activity, and over short periods (seconds to minutes) in response to acute changes in synaptic activity, as seen during long-term potentiation (LTP) or long-term depression (LTD). Modulation of AMPAR function occurs through two distinct mechanisms: modulation of ion-channel properties and regulation of

synaptic targeting of the receptors. Both processes are regulated by the phosphorylation of AMPARs (Bettler and Mulle, 1995; Dingledine 1999; Malinow and Malenka, 2002; Song and Huganir, 2002; Bredt and Nicoll, 2003; Debanne et al., 2003; Esteban, 2003; Malenka, 2003; Riedel et al., 2003; Schenk and Matteoli, 2004; Mayer, 2005; Wang et al., 2005; Groc and Choquet, 2006; Nicoll et al., 2006; Sprengel, 2006; Derkach et al., 2007; Greger et al., 2007; Greger and Esteban, 2007; Kessels and Malinow, 2009; Santos et al., 2009).

1.1.3 Kainate receptors

Kainate receptors (KARs) play important roles in regulating synaptic transmission and neuronal excitability. KARs are localized in postsynapse and presynapse. Postsynaptic KARs mediate synaptic currents of small amplitude and slow rise and decay kinetics compared with AMPAR-mediated currents. The ion channel formed by kainate receptors is permeable to sodium and potassium ions. Presynaptic KARs can modulate neurotransmitter release such as GABA and glutamate. Therefore presynaptic KARs can facilitate presynaptic forms of short and long term synaptic plasticity. KARs are tetrameric receptors composed of various combinations of five subunits GluR5, GluR6, GluR7, KA1 and KA2. GluR5, GluR6 and GluR7 can form functional homomeric and heteromeric receptor channels and have multiple isoforms derived from alternative splicing and RNA editing. GluR5 has two N-terminal splice variants (GluR5-1 and GluR5-2) and three C-terminal splice variants in the cytoplasmic domain (GluR5a, GluR5b and GluR5c). GluR6 has two C-terminal splice variants (GluR6a and GluR6b). GluR7 has two C-terminal splice variants (GluR7a and GluR7b). KA1 and KA2 can only form functional receptors by combining with one of the GluR5-7 subunits. No splice variants for KA1 and KA2 have been found (Bettler and Mulle, 1995; Bleakman, 1999; Chittajallu et al., 1999; Dingledine 1999; Kullmann, 2001; Kamiya, 2002; Debanne et al., 2003; Huettner, 2003; Lerma, 2003, 2006; Riedel et al., 2003; Isaac et al., 2004; Jaskolski, 2005; Mayer, 2005; Pinheiro and Mulle, 2006, 2008; Coussen, 2009).

1.2 Metabotropic glutamate receptors

Metabotropic glutamate receptors (mGluRs) are a family of G-protein-coupled receptors and have been shown to play an important role in synaptic plasticity, learning and memory, neuronal development and neurodegeneration (Nakanishi, 1994; Pin and Bockaert, 1995; Pin and Duvoisin, 1995; Conn and Pin, 1997; Anwyl, 1999; Bordi and Ugolini, 1999; De Blasi et al., 2001; Hermans and Challiss, 2001; Dale et al., 2002; Gerber, 2002; Valenti et al., 2002; Conn, 2003; Conn et al., 2005; Kew and Kemp, 2005; Dhami and Ferguson, 2006; Ferraguti and Shigemoto, 2006; Gerber et al., 2007).

1.2.1 Classification

Eight distinct mammalian mGluRs have been reported and have been divided into three subgroups based on sequence similarities, pharmacological agonists and antagonists, and signal transduction pathways (Nakanishi, 1994; Pin & Bockaert, 1995; Pin & Duvoisin, 1995; Conn & Pin, 1997; Dale et al., 2002). They are Group I (mGluR1 and mGluR5), Group II (mGluR2 and mGluR3) and Group III (mGluR4, mGluR6, mGluR7, and mGluR8) (Fig. 1.3). Splice variants have been reported for mGluR1, mGluR3, mGluR5, mGluR6, mGluR7 and mGluR8 (Fig. 1.3; Pin and Duvoisin 1995; Kew and Kemp, 2005; Ferraguti and Shigemoto, 2006).

Group I mGluRs, which include mGluR1 and mGluR5, are selectively activated by 3,5-dihydroxyphenylglycine (3,5-DHPG). Via G-protein-dependent pathways, group I mGluRs are positively coupled to the phospholipase C (PLC), inositol-1,4,5-trisphosphate (IP₃) formation, activation of protein kinase C (PKC) and release of calcium from intracellular stores (Abdul-Ghani et al., 1996; Conn and Pin, 1997; Bordi and Ugolini, 1999; Hermans and Challiss, 2001; Dhami and Ferguson, 2006). Group I mGluRs can also activate intracellular signals independent of G proteins, via tyrosine kinase Src (Heuss et al., 1999; Benquet




Figure 1.3 Schematic drawing of alternative splice variants of metabotropic glutamate receptors. The different colors in the C-terminus indicate the changes in amino acid sequence between isoforms attributable to alternative splicing. TMD, transmembrane domain. Taken from Ferraguti and Shigemoto, 2006.

et al., 2002; Gerber, 2002; Krause et al., 2002; Gee et al., 2003; Gee and Lacaille, 2004; Gerber et al., 2007).

Group II mGluRs, which include mGluR2 and mGluR3, are selectively activated by DCG-IV or LY379268 and are negatively coupled to adenylyl cyclase. Group II mGluRs negatively modulate glutamate and GABA neurotransmission (Anwyl, 1999; Schoepp, 2001; Kew and Kemp, 2005; Dhami and Ferguson, 2006; Ferraguti and Shigemoto, 2006).

Group III mGluRs, which include mGluR4, mGluR6, mGluR7, and mGluR8, are

selectively activated by L-(+)-2-amino-4-phosphonobutyrate (L-AP4) and are negatively coupled to adenylyl cyclase. Group III mGluRs negatively modulate glutamate and GABA neurotransmission (Anwyl, 1999; Schoepp, 2001; Kew and Kemp, 2005; Dhami and Ferguson, 2006; Ferraguti and Shigemoto, 2006).

1.2.2 Structure




Figure 1.4 Schematic drawing of the structure of an mGluR. Taken from Kew and Kemp, 2005.

The topology of mGluRs is shown in Fig. 1.4, which includes an N-terminal extracellular domain, three extracellular and three intracellular loops and a cytoplasmic C-terminal tail, separated by seven transmembrane domains. The large N-terminal extracellular domain contains the glutamate-binding site. The transmembrane heptahelical domain contains the binding sites for positive and negative allosteric modulators. The intracellular C-terminal domain interacts with cytoskeletal, scaffolding and signaling proteins (e.g., G-proteins) and with integral membrane receptors. These proteins are responsible for the correct targeting of mGluRs to specific sub-cellular compartments, coupling to effector proteins,




Figure 1.5 Binding domains of mGluRs-interacting proteins in the C-terminus of mGluRs. *Abbreviations:* CaM, Calmodulin; Siah-1A, seven in absentia homolog-1A; GRIP, glutamate receptor interacting protein; PICK1, protein interacting with C kinase 1. Taken from Fagni et al. 2004.

modulation of intracellular signalling and the recycling of mGluRs (Fig. 1.5) (De Blasi et al., 2001; De Blasi et al., 2001; Hermans and Challiss, 2001; Jingami et al., 2003; Fagni et al. 2004; Kew and Kemp, 2005; Ferraguti and Shigemoto, 2006).

1.2.3 Function

mGluRs have been implicated in diverse functions in the mammalian CNS (Pin and Duvoisin, 1995; Conn and Pin, 1997; Anwyl, 1999; Conn, 2003; Kew and Kemp, 2005; Dhami and Ferguson, 2006; Ferraguti and Shigemoto, 2006). mGluRs mediate slow excitatory (Charpak and Gähwiler, 1991; Glaum and Miller 1992; McCormick and von Krosigk 1992; Eaton et al. 1993; Congar et al., 1997; Gee et al., 2003; Wu et al. 2004) and inhibitory (Fiorillo and Williams 1998) responses. mGluRs regulate calcium channels (Swartz and Bean 1992; Sahara and Westbrook 1993; Chavis et al. 1994), potassium channels (Charpak et al. 1990; Guérineau et al., 1994; Shirasaki et al. 1994; Gereau and Conn 1995b;

Netzeband et al. 1997) and non-selective cation channels (Guérineau et al. 1995; Congar et al. 1997). mGluRs regulate the trafficking of ionotropic glutamate receptors (Lan et al. 2001; Snyder et al. 2001) and modify NMDAR-mediated synaptic transmission (Awad et al. 2000; Aniksztejn et al., 1991, 1992; Bleakman et al., 1992; Harvey and Collingridge, 1993; Pisani et al., 1997, 2001; Ugolini et al., 1997, 1999; Yu et al., 1997; Contractor et al., 1998; Wang et al., 1998; Zhong et al., 2000; Benquet et al., 2002; Zho et al., 2002; Grishin et al., 2004; Guo et al. 2004; Harney et al. 2006; Ireland and Abraham, 2009). mGluRs inhibit (Baskys and Malenka 1991; Desai and Conn 1991; Schrader and Tasker 1997; Wittmann et al. 2001; Xi et al. 2003; Acuna-Goycolea et al. 2004; Chu and Moenter 2005) and facilitate transmitter release (Rodriguez-Moreno et al. 1998; Morsette et al. 2001; Herrero et al. 1992). Activation of mGluRs can induce LTP (Bortolotto and Collingridge 1993; O'Connor et al. 1995; Manahan-Vaughan 1997; Raymond et al. 2000; Chiamulera et al., 2001; Miura et al. 2002; Gubellini et al., 2003; Sourdet et al., 2003; Topolnik et al., 2006; Anwyl, 2009) and LTD (Linden et al. 1991; Stanton et al., 1991; Bashir et al., 1993b; Kato 1993; Bolshakov and Siegelbaum 1994; Conquet et al. 1994; Shigemoto et al. 1994; Kemp and Bashir, 1997, 1999; Oliet et al., 1997; Palmer et al., 1997; Nicoll et al., 1998; Fitzjohn et al., 1999; Huber et al., 2000, 2001; Lin et al. 2000; Gubellini et al., 2001; Xiao et al., 2001; Daw et al., 2002; Feinmark et al., 2003; Otani et al. 2002; Gallagher et al., 2004; Moulton et al., 2006; Grueter et al., 2007; Cho et al., 2008; Jo et al., 2008; Neyman and Manahan-Vaughan, 2008; Wang et al., 2008; Waung et al., 2008; Holbro et al., 2009). mGluRs play an important role in memory formation (Aiba et al. 1994; Packard et al. 2001) and the regulation of neuronal development (Pin and Duvoisin 1995; Hensch and Stryker 1996; Conn and Pin 1997; Kano et al. 1997; Plenz and Kitai 1998; Catania et al. 2001; Hannan et al. 2001). mGluRs are involved in various diseases such as anxiety and stress disorders (Linden et al. 2002; Swanson et al. 2005), epilepsy (Sansig et al. 2001), excitotoxic neuronal death (Bruno et al. 2000), hypoxic brain damage (Poli et al. 2003) and fragile X mental retardation (Bear et al. 2004).

1.3 Group I metabotropic glutamate receptors

1.3.1 Distribution of group I mGluRs in the central nervous system

mGluR1 has been found in hippocampal neurons, in Purkinje cells of the cerebellar cortex, in mitral/tufted cells of the olfactory bulb, as well as in the lateral septum, globus pallidus, entopeduncular nucleus, ventral pallidum, magnocellular preoptic nucleus and thalamus (Martin et al. 1992; Shigemoto et al. 1992; Hubert et al. 2001).

Differential mRNA expression of mGluR1 splice variants has been observed (Berthele et al. 1998, 1999). Strong expression of the mGlu1 α splice variant has been found in Purkinje cells, hippocampal interneurons, mitral and tufted cells, thalamic neurons and in the substantia nigra (Martin et al. 1992; Baude et al. 1993; Fotuhi et al. 1993; Ferraguti et al. 2004). The mGlu1 β splice variant is intensively expressed in CA3 hippocampal pyramidal neurons, granule cells of the dentate gyrus and lateral hypothalamus (Ferraguti et al. 1998; Mateos et al. 1998).

Protein levels of mGluR1 α are found to increase progressively during pre- and post-natal development (Catania et al. 1994; van den Pol et al. 1994; Lopez-Bendito et al. 2002; Shigemoto et al. 1992).

In contrast, the expression of mGluR5 is low in the cerebellum (Neki et al. 1996; Negyessy et al. 1997). High expression has been found in the cerebral cortex, hippocampus, subiculum, olfactory bulb, anterior olfactory nucleus, olfactory tubercle, striatum, nucleus accumbens and lateral septal nucleus (Abe et al. 1992; Shigemoto et al. 1993; Romano et al. 1995; Bordi and Ugolini, 1999). In the hippocampus, mGluR5 is mainly expressed in interneurons as well as in dendritic fields of pyramidal and granule cells (Kerner et al. 1997; Shigemoto et al. 1997).

mGluR5 increases perinatally, with a peak around the second postnatal week, and decreases thereafter (Catania et al. 1994; Romano et al. 1996; Lopez-Bendito et al. 2002). Changes in mGluR5 expression have been associated with a gradual decrease in mGluR5a and an increase in mGluR5b. mGluR5a is found most abundantly in the young rat, whereas mGluR5b predominates in the adult rat (Joly et al. 1995; Minakami et al. 1995; Romano et al. 1996).

1.3.2 Subcellular localization of group I mGluRs

At the electron-microscopic level, although Group I mGluRs can be located at both presynaptic and postsynaptic sites in the brain (Gereau and Conn, 1995a; Manzoni and Bockaert, 1995; Romano et al. 1995; Jia et al. 1999; Simkus and Stricker, 2002; White et al., 2003; Park et al., 2004; Musante et al., 2008), they are mostly observed at the periphery of the postsynaptic densities of asymmetrical glutamatergic synapses (Baude et al., 1993; Shigemoto et al., 1993, 1997; Nusser et al. 1994; Lujan et al., 1996, 1997; Vidnyánszky et al. 1996; Liu et al. 1998).

mGluR1 has been detected mostly postsynaptically in neurons in the hippocampus (Baude et al. 1993; Lujan et al. 1996, 1997; Hanson and Smith 1999), cerebral cortex (Ong et al. 1998), striatum (Hanson and Smith 1999), thalamus (Martin et al. 1992; Godwin et al. 1996; Liu et al. 1998), hypothalamus (van den Pol et al. 1994), cerebellar cortex (Martin et al. 1992; Baude et al. 1993; Gorcs et al. 1993; Nusser et al. 1994; Lujan et al. 1996, 1997; Jaarsma et al. 1998; Mateos et al. 2000) and dorsal cochlear nucleus (Petrálie et al. 1996; Jaarsma et al. 1998). At the head of Purkinje cell spines in the rat cerebellar cortex, about half of the immunogold particles for mGluR1 α are localized perisynaptically, i.e. within a 60-nm annulus surrounding the edge of synapses, whereas the remaining particles are localized extrasynaptically at more distant positions (Lujan et al. 1997).

mGluR5 has been detected mainly in somatic and dendritic domains in the hippocampus (Lujan et al. 1996, 1997; Hanson and Smith 1999), basal ganglia (Shigemoto et al. 1993; Hanson and Smith 1999), thalamus (Godwin et al. 1996; Liu et al. 1998), hypothalamus (Romano et al. 1995; van den Pol et al. 1995), cerebellar cortex (Négyessy et al. 1997), dorsal cochlear nucleus (Petrálie et al. 1996) and dorsal horn of the spinal cord (Vidnyánszky et al. 1994; Jia et al. 1999). In a few studies, mGluR5 immunoreactivity has been reported not only in somatodendritic domains of neurons, but also in axons (Romano et al. 1995) or vesicle-containing profiles (Jia et al. 1999). At the head of CA1 pyramidal neuron spines, about one fourth of the immunogold particles for mGluR5 are observed at the periphery of the postsynaptic density (PSD) of asymmetrical synapses (Lujan et al. 1996, 1997).

In the developing nervous system, mGluR5 is preferentially found in unmyelinated axons and pyramidal cell dendrites in the first two postnatal weeks. As development proceeds, the number of mGluR5-immunolabelled unmyelinated axons is significantly reduced and the mGluR5 becomes more widely distributed on dendritic spines (Liu et al. 1998; Lopez-Bendito et al. 2002; Hubert and Smith 2004).

1.3.3 Function of group I mGluRs

Group I mGluRs have been shown to play an important role in neuronal development (Catania et al., 1991; Plenz & Kitai, 1998; Flint et al., 1999; Hannan et al., 2001), synaptic plasticity (Zhong et al., 2000; Wu et al., 2001; Gubellini et al., 2003), as well as in several neurological and neurodegenerative disorders (Kingston et al., 1999; Bruno et al., 2001), such as chronic pain (Bhave et al., 2001; Karim et al., 2001; Zhou et al., 2001), epilepsy (Chapman et al., 2000), multiple sclerosis (Geurts et al., 2003), amyotrophic lateral sclerosis (Laslo et al., 2001), Huntington's disease (Calabresi et al., 1999) and Parkinson's disease (Awad et al., 2000).

The activation of group I mGluRs has also been shown to play an important role in long- and short-term synaptic plasticity. Two forms of long-term synaptic plasticity, LTP and LTD, are observed in several brain regions and have been proposed to represent the cellular mechanisms for learning and behaviour. The activation of group I mGluRs is required for the induction of LTP (Bashir et al., 1993a; Aiba et al., 1994; Conquet et al., 1994; O'Connor et al., 1995; Lu et al., 1997; Raymond et al., 2000; Chiamulera et al., 2001; Gubellini et al., 2003; Sourdet et al., 2003; Topolnik et al., 2006; Anwyl, 2009) and LTD (Linden et al., 1991; Bashir et al., 1993b; Kato 1993; Bolshakov and Siegelbaum 1994; Conquet et al., 1994; Shigemoto et al. 1994; Kemp and Bashir, 1997, 1999; Oliet et al., 1997; Palmer et al., 1997; Nicoll et al., 1998; Fitzjohn et al., 1999; Huber et al., 2000, 2001; Gubellini et al., 2001; Xiao et al., 2001; Daw et al., 2002; Feinmark et al., 2003; Gallagher et al., 2004; Moulton et al., 2006; Grueter et al., 2007; Cho et al., 2008; Jo et al., 2008; Neyman and Manahan-Vaughan, 2008; Wang et al., 2008; Waung et al., 2008; Holbro et al., 2009). In the hippocampus, mGluR1-deficient mice exhibit either normal or only slightly reduced CA1 and dentate LTP, but impaired CA3 mossy fiber LTP (Aiba et al., 1994; Conquet et al., 1994). Conversely, mGluR5-deficient mice display a reduction in CA1 hippocampal LTP, but demonstrate normal CA3 mossy fiber LTP and have a reduced performance in spatial learning tasks (Lu et al., 1997).

It was reported that group I mGluRs modulate NMDA receptor responses in CNS (Awad et al. 2000; Aniksztejn et al., 1991, 1992; Bleakman et al., 1992; Harvey and Collingridge, 1993; Pisani et al., 1997, 2001; Ugolini et al., 1997, 1999; Yu et al., 1997; Contractor et al., 1998; Wang et al., 1998; Zhong et al., 2000; Lan et al., 2001; Snyder et al., 2001; Benquet et al., 2002; Zho et al., 2002; Grishin et al., 2004; Guo et al., 2004; Harney et al., 2006; Ireland and Abraham, 2009), which may represent an additional mechanism through which NMDA-dependent synaptic plasticity can be modulated.

1.4 Long-term potentiation and long-term depression

1.4.1 Long-term potentiation

Long-term potentiation (LTP) of the chemical synaptic transmission is the enhancement of signal transmission between two nerve cells which lasts for an extended period of time (minutes to hours *in vitro* and hours to days and months *in vivo*). LTP is widely considered as the major cellular basis of learning and memory (Teyler and DiScenna, 1987; Bliss and Collingridge, 1993; McNaughton, 1993; Martinez and Derrick, 1996; Redman, 1996; Malenka and Nicoll, 1999; Debanne et al., 2003; Malenka and Bear, 2004; Cooke and Bliss, 2006; Kullmann and Lamsa, 2007).




Figure 1.6 NMDAR-dependent LTP. Depolarization of the postsynaptic cell relieves Mg^{2+} block of the NMDAR, allowing Na^+ and Ca^{2+} to flow into the dendritic spine through the NMDAR. The resultant rise in Ca^{2+} within the dendritic spine is the critical trigger for LTP. Taken from Malenka and Nicoll, 1999.

LTP requires coincident activity of presynaptic and postsynaptic elements, resulting in a facilitation of transmission. Homosynaptic LTP can be induced experimentally by applying a sequence of short, high-frequency stimulations to the presynaptic cell. Some types of LTP depend on the activation of postsynaptic NMDARs (Fig. 1.6), which result in calcium influx in the postsynaptic cell and the activation of protein kinases, i.e., protein kinase C (PKC) and calcium/calmodulin-dependent protein kinase II (CaMKII) (Teyler and DiScenna, 1987; Anwyl, 1989; Madison et al., 1991; Bashir and Collingridge, 1992; Bliss and Collingridge, 1993; Malenka, 1994; Kullmann and Siegelbaum, 1995; Malenka and Nicoll, 1999; Matynia et al., 2002; Malenka and Bear, 2004; Kullmann and Lamsa, 2007). Other types are NMDAR-independent (Johnston et al., 1992; Bear and Malenka, 1994; Bortolotto et al., 1999; Malenka and Bear, 2004; Kullmann and Lamsa, 2007). However, whether the induction of LTP relies exclusively on a postsynaptic process (Matsuzaki et al., 2004; Kerchner and Nicoll, 2008), or rather involves a presynaptic reduction in neurotransmitter release probability (Bayazitov et al., 2007; Ahmed and Siegelbaum, 2009; Enoki et al., 2009), remains under debate (Madison et al., 1991; Malenka, 1994; Isaac et al., 1996; Lisman, 2009).

1.4.2 Long-term depression

The opposite phenomenon of LTP, called long-term depression (LTD), is the weakening of a neuronal synapse that lasts for an extended period of time (minutes to hours *in vitro* and hours to days and months *in vivo*). Cerebellar LTD has been hypothesized to be important for motor learning. Hippocampal LTD may mediate learning, forgetting or behavioural extinction, e.g., the loss of responsiveness to previously effective stimuli. Alternatively, LTD may serve as a homeostatic mechanism to ensure that CNS synapses are not saturated by learning (Ito, 1989; Linden, 1994; Bear and Malenka, 1994; Malenka, 1994; Linden and Connor, 1995; Bear and Abraham, 1996; Debanne and Thompson, 1996; Thiels et al., 1996; Kemp and Bashir, 2001; Debanne et al., 2003; Anwyl,




Figure 1.7 Types of LTD. Vertical lines represent stimulating pulses. “—” sign shows the location of LTD. Abbreviations: PN: Purkinje neuron; CF: climbing fiber; PF: parallel fiber. Adapted from Linden, 1994

2006; Cooke and Bliss, 2006; Kullmann and Lamsa, 2007).

There are several types of LTD (Fig. 1.7). In heterosynaptic LTD, inducing stimuli delivered to one pathway result in a depression in the pathway that did not receive stimuli. In homosynaptic LTD, depression is confined to the stimulated pathway. Induction of associative LTD and cerebellar LTD require the activation of two pathways (Linden, 1994; Linden and Connor, 1995; Debanne and Thompson, 1996; Kemp and Bashir, 2001).

The standard protocol for inducing homosynaptic LTD is a long train of low-frequency (1 Hz) stimulation (Dudek and Bear, 1992; Mulkey and Malenka, 1992). Coincidence of pre- and post-synaptic action potentials can induce LTD as well

(Markram et al., 1997). NMDAR-dependent (Bear and Malenka, 1994; Linden and Connor, 1995; Bear and Abraham, 1996; Thiels et al., 1996; Kemp and Bashir, 2001; Malenka and Bear, 2004) and mGluR-dependent (Linden and Connor, 1995; Bear and Abraham, 1996; Bortolotto et al., 1999; Kemp and Bashir, 2001; Malenka and Bear, 2004; Anwyl, 2006) homosynaptic LTD have been characterized.

1.5 Oscillations

1.5.1 Function of oscillations

Network oscillations *in vivo* have been proposed to be important in sensory processing (Laurent and Davidowitz, 1994), in sensory and perceptual binding (Singer and Gray, 1995; Miltner et al., 1999), in motor programming (Murthy and Fetz, 1996), in associative learning (Rodriguez et al., 1999; Buzsáki, 2002), and in epileptogenesis (Grenier et al., 2001, 2003).

1.5.2 Oscillations in hippocampus

In the behaving animals, synchronous oscillations of the membrane potential of large neuronal populations can occur at various frequencies in the hippocampus (Table 1.1).

1.5.3 Methods of inducing oscillation *in vitro*

A variety of population oscillations can be induced in hippocampal slices either by activation of metabotropic glutamate receptors, kainate receptors or muscarinic receptors with drugs or electrical stimulation; or by manipulation of the ionic

Oscillation type	Conditions	Chemical synapses	GJ	Pyramidal cell somata	Interneurons
ING	mGluR + iGluR block	Yes	Yes	—	Each wave
	[K ⁺] puff + iGluR block	Yes	Yes	—	N.T.
tetanic γ	Elect. stim.	Yes	No?	Variable: can be most beats [pyr. cells and interneurons in phase]	Most/all beats; doubles if two-site
tetanic β	Elect stim. (two-site, strong)	Yes	No?	Skip beats	γ frequency
“pure” VFO	spontaneous or low [Ca ²⁺]	No	Yes	Some waves; spikelets	—
persistent γ	kainate, mAChR, mGluR	Yes	Yes	Rare firing	Fire 1/2 to most waves (axo-axonic even more)
				[pyr. cells slightly lead interneurons]	
puff γ	Local hypertonic [K ⁺]	Yes	Yes	Rare firing	N.T.
theta (CA1)	mGluR + AMPA block	Yes	Yes	Rare firing	S. oriens interneurons (O/LM) fire each wave
theta (CA3)	mAChR	Yes (not GABA _A)	N.T.	May fire little, or may burst	Probably not necessary

Table 1.1 Properties of *in vitro* hippocampal oscillations. Abbreviations: N.T., not tested; ING: interneuron network gamma; VFO, very fast oscillation (>70 Hz); gamma/beta (15–70 Hz); theta rhythm (4-15 Hz). Taken from Traub et al., 2004

environment (for example, removing of calcium ions); brief oscillations can even occur spontaneously (Traub et al., 2004).

1.5.4 Mechanisms of the generation of theta rhythm

Theta rhythm (4-15 Hz) (Bland 1986; Lopes da Silva et al. 1990; Stewart and Fox

1990; Vinogradova 1995; Vertes and Kocsis 1997; Kamondi et al., 1998; Harris et al., 2001; Buzsáki, 2002) is present mainly during locomotion and other voluntary movements (Grastyán et al., 1959; Vanderwolf, 1969) and rapid eye movement (REM) sleep (Jouvet, 1969). Theta waves have been also assumed to code mnemonic processes (Lisman and Idiart 1995; Raghavachari et al. 2001). Loss of hippocampal theta rhythm results in spatial memory deficits in the rat (Winson 1978).

Mechanisms of the generation of theta rhythm in the hippocampus are not completely understood yet. However, it is clear that the reciprocal connection between the medial septum-diagonal band of Broca (MS-DBB) and the hippocampus plays an essential role (Green and Arduni, 1954; Petsche et al., 1962; Lawson and Bland, 1993; Sik et al., 1994; Markowska et al., 1995; King et al., 1998; Dragoi et al., 1999; Buzsáki 2002; Gulyas et al., 2003). Hippocampal interneurons represent the only output from the hippocampus to the MS-DBB (Toth et al., 1993) and are required for the induction and propagation of theta rhythm (Buzsáki et al., 1983; Sik et al., 1994; Freund and Antal, 1988; Buzsáki and Chrobak, 1995; Cobb et al., 1995; Dragoi et al., 1999; Buzsáki 2002; Gulyas et al., 2003; Mann and Paulsen, 2007). One hippocampal interneuron innervates more than 1,000 pyramidal cells (Halasy et al., 1996). Thus, they are in a key position to synchronize the activity of pyramidal cells.

Chapter 2

Group I metabotropic glutamate receptors in CA1 pyramidal cells are not tonically activated by ambient glutamate in hippocampal slice cultures

2.1 Abstract

Microdialysis experiments report a background level of extracellular glutamate concentration of approximately 2 μM in the brain. The EC_{50} of glutamate for group I mGluRs is $\sim 10 \mu\text{M}$. Therefore the group I mGluRs may be tonically activated by the background level of glutamate present in the extracellular space. Here we recorded in CA1 pyramidal neurons and found that the ambient glutamate concentration in the extracellular space is insufficient to tonically activate group I mGluRs in hippocampal organotypic slice cultures.

2.2 Introduction

The concentration of neurotransmitters in the extracellular space of the central nervous system is determined by a balance between release, degradation and uptake mechanisms. Glutamate is released during fast synaptic transmission and its concentration reaches $\sim 1 \text{ mM}$ within the synaptic cleft (Clements, 1996). Then glutamate concentration decreases rapidly due to diffusion and uptake by glutamate transporter (Bergles et al. 1999; Attwell and Gibb, 2005). If the release rate is low, in principle, the co-transport of 3 Na^+ and 1 H^+ and the counter-transport of 1 K^+ can lower the extracellular glutamate concentration to $\sim 2 \text{ nM}$ (Zerangue and Kavanaugh, 1996; Levy et al., 1998). However, microdialysis experiments report extracellular glutamate concentrations of 0.5-7 μM (typically 2 μM) in the brain *in vivo* and in perfusate from unstimulated brain slices (Johnson 1978; Benveniste et al., 1984; Phillis et al., 1994; Wahl et al., 1994; Bianchi et al. 1999; Meldrum 2000), which are higher than these theoretical values. The EC_{50} of glutamate for NMDA receptors is $\sim 2 \mu\text{M}$ (Patneau and Mayer, 1990; Hollmann and Heinemann 1994; Meldrum 2000; Nahum-Levy et al., 2001, 2002). Accordingly, tonic activation of NMDA receptors in pyramidal neurons was

observed in acute brain slices (Sah et al., 1989; Angulo et al., 2004; Cavelier and Attwell, 2005; Cavelier et al., 2005; Meur et al., 2007).

The EC₅₀ of glutamate for group I mGluRs is ~10 μ M (Hollmann and Heinemann 1994; Pin and Duvoisin 1995; Conn and Pin, 1997). Therefore the group I mGluRs may be tonically activated by the background level of glutamate present in the extracellular space. Group I mGluRs modulate NMDA receptor-mediated responses in CA1 pyramidal cells (Aniksztejn et al., 1991, 1992; Harvey and Collingridge, 1993; Contractor et al., 1998; Snyder et al., 2001; Benquet et al., 2002; Zho et al., 2002; Grishin et al., 2004; Ireland and Abraham, 2009). Here we used the activity of NMDA receptors as a sensor to check whether the ambient glutamate concentration in the extracellular space is sufficient to tonically activate group I mGluRs in hippocampal slice cultures.

2.3 Materials and Methods

2.3.1 Slice cultures and electrophysiology

Hippocampal organotypic slice cultures were prepared from 6-day-old Wister rats using the roller-tube techniques, as described previously (Gähwiler et al., 1998). After 3-4 weeks *in vitro* the slice cultures were transferred to a recording chamber with a volume of 1 ml on an upright microscope (Axioskop FS, Zeiss). Slices were superfused at a rate of 1.5 ml/min with artificial cerebrospinal fluid (ACSF, in mM) 137 NaCl, 2.7 KCl, 11.6 NaHCO₃, 0.4 NaH₂PO₄, 2.0 MgCl₂, 2.8 CaCl₂, 5.6 D-glucose and 10 mg/L phenol red (pH 7.4, ~305 mOsm, 33 °C, saturated with 95% O₂ and 5% CO₂).

Whole-cell voltage-clamp recordings were obtained from hippocampal CA1 pyramidal cells with an Axopatch 200B amplifier (Molecular Devices). Recording

pipettes (3-5 M Ω) were filled with (in mM) 120 K-gluconate, 10 L-glutamic acid, 5 KCl, 10 Hepes, 10 EGTA, 2 MgATP, 5 creatine phosphate (CrP), 0.4 NaGTP, 0.07 CaCl₂ (pH 7.2, ~290 mOsm). Membrane potentials were corrected for liquid junction potentials. Series resistance (5-15 M Ω) and input resistance were monitored regularly. Currents were filtered at 5 kHz and analyzed off-line (pClamp 9, Molecular Devices). All numerical data are presented as the mean \pm SEM. Significance was tested using two-tailed t-test.

2.3.2 Induction of NMDA currents

CA1 pyramidal cells were voltage-clamped at -50 mV. In the presence of 1 μ M TTX (sodium channel blocker), 100 μ M picrotoxin (GABA_A receptor antagonist) and 10 μ M NBQX (AMPA/kainate receptors antagonist), NMDA currents were induced by applying brief pressure pulses (1 bar, ~200 msec) at 40-second intervals to a micropipette filled with NMDA (100 μ M) and placed ~100 μ m from the soma of the recorded cell.

2.3.3 Drugs

NBQX (2,3-dioxo-6-nitro-1,2,3,4-tetrahydrobenzo[f]quinoxaline-7-sulfonamide) and (S)-MCPG ((S)- α -methyl-4-carboxyphenylglycine) were purchased from Ascent Scientific (Bristol, UK). All other chemicals were purchased from Sigma/Fluka.

2.4 Results

2.4.1 Group I mGluRs in CA1 pyramidal cells are not tonically activated by ambient glutamate in hippocampal slice cultures

CA1 pyramidal cells were voltage-clamped at -50 mV. In the presence of 1 μ M TTX (sodium channel blocker), 100 μ M picrotoxin (GABA_A receptor antagonist) and 10 μ M NBQX (AMPA/kainate receptors antagonist), NMDA currents were induced by applying brief pressure pulses (1 bar, ~200 msec) at 40-second intervals to a micropipette filled with NMDA (100 μ M) and placed ~100 μ m from the soma of the recorded cell (Fig. 2.1A). Application of the group I mGluRs antagonist (S)-MCPG did not modulate NMDA receptor-mediated responses in CA1 pyramidal cells (500 μ M; $102 \pm 3\%$ of baseline, $n = 12$, $p = 0.8$; Fig. 2.1B). In other words, NMDA receptors were not tonically modulated by group I mGluRs. MCPG did not change the holding current of pyramidal cells voltage-clamped at -50 mV ($n = 12$, $p = 0.5$), indicating that there was no tonic current mediated by group I mGluRs. Take together, we conclude that group I mGluRs in CA1 pyramidal cells are not tonically activated by the background level of glutamate present in the extracellular space.




Figure 2.1 Group I metabotropic glutamate receptors in CA1 pyramidal cells are not tonically activated by ambient glutamate in hippocampal slice cultures. (A) Schematic drawing of induction of NMDA currents. (B) Application of 500 μ M (S)-MCPG did not modulate NMDA receptor-mediated responses in CA1 pyramidal cells ($n = 12$).

2.5 Discussion

This study shows that there is no tonic activation of group I mGluRs in CA1 pyramidal cells in hippocampal organotypic slice cultures.

Microdialysis experiments report a background level of extracellular glutamate concentration of 0.5-7 μM (typically 2 μM) in the brain *in vivo* and in perfusate from unstimulated brain slices (Johnson 1978; Benveniste et al., 1984; Phillis et al., 1994; Wahl et al., 1994; Bianchi et al. 1999; Meldrum 2000). Our experiments are performed in hippocampal organotypic slice cultures, in which the background level of extracellular glutamate may be lower than in the brain *in vivo* or in acute brain slices. Therefore a tonic activation of group I mGluRs cannot be observed in hippocampal organotypic slice cultures.

A recent study shows that the ambient concentration of glutamate in acute hippocampal slices is ~25 nM (Herman and Jahr, 2007), 100-fold lower than former reports (Johnson 1978; Benveniste et al., 1984; Phillis et al., 1994; Wahl et al., 1994; Bianchi et al. 1999; Meldrum 2000) and is inconsistent with the reports of tonic activation of NMDA receptors (Sah et al., 1989; Angulo et al., 2004; Cavelier and Attwell, 2005; Cavelier et al., 2005; Meur et al., 2007). However, it may explain why tonic activation of group I mGluRs is not observed in my study.

Consistent with the above result, in chapter 3, we find that multiple Shaffer collateral inputs are required to activate group I mGluRs in CA1 pyramidal cells.

Chapter 3

Activation conditions for the induction of metabotropic glutamate receptor-dependent LTD in hippocampal CA1 pyramidal cells

3.1 Abstract

Two forms of homosynaptic long-term depression (LTD) are distinguished in hippocampal CA1 pyramidal cells, one which is NMDA receptor-dependent and the other metabotropic glutamate receptor (mGluR)-dependent. Although the molecular processes involved in mGluR-LTD are well characterized, the conditions of circuit activation required for its induction remain unclear. We show that mGluR-LTD cannot be induced in synaptically coupled CA3-CA1 pyramidal cell pairs. Experiments to address the underlying mechanisms indicate that even when glutamate transporters are blocked, one presynaptic cell releases insufficient glutamate to evoke an mGluR-mediated current in a connected CA1 cell. These findings imply that extrasynaptic diffusion is not a limiting factor, and are consistent with a sparse distribution of functional mGluRs in the dendritic tree of pyramidal cells. Thus, the discharge of multiple Schaffer collaterals to a targeted cell is necessary for mGluR-LTD. Our experiments indicate that approximately eight CA3 inputs to a CA1 pyramidal cell must be activated to induce mGluR-LTD.

3.2 Introduction

Homosynaptic LTD in the hippocampus is well characterized as an NMDA receptor-dependent form of synaptic plasticity (NMDAR-LTD), as first described at the Schaffer collateral input to CA1 pyramidal cells (Dudek and Bear, 1992; Mulkey and Malenka, 1992). An additional form of LTD at Schaffer collateral synapses onto CA1 pyramidal cells that requires activation of mGluRs (mGluR-LTD) was also reported (Stanton et al., 1991; Bolshakov and Siegelbaum, 1994;

Oliet et al., 1997). Significant progress has been made in understanding the induction mechanisms and intracellular transduction pathways involved in each form (Kemp and Bashir, 2001; Anwyl, 2006). Both appear to be induced postsynaptically (Oliet et al., 1997), but whether their expression relies exclusively on postsynaptic processes (Snyder et al., 2001; Xiao et al., 2001; Zhou et al., 2004; Zhang et al., 2006; Brager and Johnston, 2007), or rather involves presynaptic reduction in neurotransmitter release probability (Bolshakov and Siegelbaum, 1994; Fitzjohn et al., 2001; Watabe et al., 2002; Enoki et al., 2009), remains under debate (Lisman, 2009). For the induction of NMDAR-LTD, it has been shown that cooperative interactions of multiple presynaptic inputs are not essential, such that appropriately timed discharge of a single CA3 pyramidal cell can lead to homosynaptic LTD in a targeted CA1 pyramidal cell (Debanne et al., 1996). A corresponding investigation of the circuit requirements for induction of mGluR-LTD is at present lacking. Here we recorded from synaptically connected pairs of CA3 and CA1 hippocampal pyramidal cells in organotypic slice cultures, and find that in contrast to NMDA-LTD, multiple Schaffer collateral input is required for the induction of mGluR-LTD in a targeted CA1 pyramidal cell.

3.3 Materials and Methods

3.3.1 Slice cultures and electrophysiology

For the preparation of hippocampal organotypic slice cultures, see chapter 2.

Recordings were obtained from hippocampal pyramidal cells with an Axopatch 200B amplifier (Molecular Devices). Both low resistance (3-5 M Ω) and high resistance (10-15 M Ω) recording pipettes were used, which were filled with (in

mM) 120 K-gluconate, 10 L-glutamic acid, 5 KCl, 10 Hepes, 1 EGTA, 2 MgATP, 5 creatine phosphate (CrP), 0.4 NaGTP, 0.07 CaCl_2 (pH 7.2, ~290 mOsm). For paired recordings, the postsynaptic pipette was filled with (in mM) 126.6 CsF, 8.4 CsCl, 10 Hepes, 10 EGTA, 2 MgATP, 5 creatine phosphate, 0.4 NaGTP (pH 7.2, ~290 mOsm). Membrane potentials were corrected for liquid junction potentials. Signals were filtered at 5 kHz and analyzed off-line (pClamp 9, Molecular Devices). All numerical data are presented as the mean \pm SEM. Significance was tested using two-tailed t-test.

3.3.2 Extracellular stimulation

For monopolar extracellular stimulation of the Schaffer collaterals, glass pipettes filled with ACSF were placed in the stratum radiatum, 100-200 μm away from the recorded cell. For the experiments depicted in supplemental Fig. 3, the monopolar stimulation pipette was positioned in CA3 stratum oriens to ensure the selective activation of Schaffer collaterals. LTD was induced with a standard low frequency stimulation (LFS) protocol (1 Hz, 900 pulses, 100 μs duration; Dudek and Bear, 1992; Manahan-Vaughan, 1997; Wang et al., 2008), or with a paired pulse low frequency stimulation (PP-LFS) protocol (1 Hz, 900 pairs of pulses, 50 ms inter-stimulus interval, 100 μs duration; Kemp and Bashir, 1997) in hippocampal CA1 pyramidal cells voltage-clamped at -70 mV in the presence of the NMDAR antagonist D-AP5 (50 μM). The stimulation intensity was 10-20 μA to induce a response of 50-70% of the maximum. To evoke a current mediated by group I mGluRs, high frequency stimulation (100 Hz, 1-4 pulses, 10-50 μA , 100 μs duration) was applied. CA1 pyramidal cells were voltage-clamped at -50 mV. AMPA, NMDA, GABA_A and GABA_B receptors were blocked by CNQX (50 μM), D-AP5 (100 μM), picrotoxin (100 μM) and CGP 62349 (5 μM), respectively.

3.3.3 Perforated patch-clamp

Recording pipettes (3-5 M Ω) were filled with (in mM) 120 K-gluconate, 10 L-glutamic acid, 5 KCl, 10 Hepes, 1 EGTA, 2 MgATP, 5 creatine phosphate, 0.4 NaGTP, 2.5 CaCl₂ (pH 7.2, ~290 mOsm). Stock solution of amphotericin B (60 mM) was prepared in DMSO and added to the pipette solution to a final concentration of 0.3 mM. A fresh solution was prepared every 1.5 h, sonicated 5-10 min and kept in the dark.

3.3.4 Drugs

CNQX (6-cyano-7-nitroquinoxaline-2,3-dione), D-AP5 (D-(-)-2-amino-5-phosphonopentanoic acid) and (S)-MCPG ((S)- α -methyl-4-carboxyphenylglycine) were purchased from Ascent Scientific (Bristol, UK). DL-TBOA (DL-threo-b-benzyloxyaspartic acid) was purchased from Tocris (Bristol, UK). CGP 62349 was kindly provided by Novartis (Basel, Switzerland). All other chemicals were purchased from Sigma/Fluka.

3.4 Results

3.4.1 A single CA3 pyramidal cell does not induce mGluR-LTD in a connected CA1 pyramidal cell

In the presence of D-AP5 (50 μ M), a standard LTD protocol evoking a low frequency train of action potentials in a CA3 pyramidal cell (1 Hz, 900 APs) failed to induce mGluR-LTD in a synaptically connected CA1 pyramidal cell in hippocampal slice cultures ($98 \pm 4\%$ of baseline, $n = 6$, $p = 0.5$, recording pipette resistance: 3-5 M Ω ; Fig. 3.1A; Fig. 3.2). Because certain forms of synaptic plasticity are disrupted by cytoplasmic washout (Kullmann and Lamsa, 2007), this experiment was repeated using high resistance recording pipettes (10-15 M Ω). Again, LTD was not observed ($98 \pm 3\%$ of baseline, $n = 3$, $p = 0.6$; Fig. 3.1A). Paired-pulse low frequency stimulation (PP-LFS) represents a more efficient protocol for the induction of mGluR-LTD (Kemp and Bashir, 1997). However, even PP-LFS (1 Hz, 900 paired APs, 50 ms inter-stimulus interval, 10-15 M Ω for the CA1 electrode) did not induce mGluR-LTD in CA3-CA1 connected cell pairs ($99 \pm 4\%$ of baseline, $n = 4$, $p = 0.6$; Fig. 3.1B). As postsynaptic mGluRs in hippocampal pyramidal cells are localized in perisynaptic and extrasynaptic zones (Lujan et al., 1996), membrane transporters may prevent glutamate from accessing these receptors. But even in the presence of TBOA (10 μ M), a glutamate transporter blocker (Jabaudon et al., 1999), LTD was not induced ($102 \pm 6\%$ of baseline, $n = 3$, $p = 0.8$; Fig. 3.1B). Taken together, these results suggest that activity in a single CA3 pyramidal cell is insufficient to induce mGluR-LTD in a targeted CA1 pyramidal cell. In contrast, when Schaffer collaterals were stimulated with an extracellular electrode (LFS, stimulation intensity: 10-20 μ A), thereby activating several CA3 pyramidal cell axons, mGluR-LTD was consistently observed ($70 \pm 3\%$ of baseline, $n = 8$, $p < 0.001$; Fig. 3.1C). Interestingly, mGluR-LTD induced by extracellular stimulation was significantly enhanced when recording with high resistance pipettes ($42 \pm 3\%$ of




Figure 3.1 Activation of a single CA3 pyramidal cell is insufficient to induce mGluR-LTD in a synaptically connected CA1 pyramidal cell. (A) In the presence of D-AP5, a low frequency train of action potentials (1 Hz, 900 APs) in a CA3 cell did not induce mGluR-LTD in a CA1 cell ($n = 6$), even when high resistance recording pipettes were used to minimize dialysis ($n = 3$). (B) A low frequency train of paired action potentials (PP-LFS) also failed to induce mGluR-LTD ($n = 4$) even when glutamate transporters were blocked with TBOA ($n = 3$). (C) mGluR-LTD can be induced in a CA1 pyramidal cell with extracellular stimulation of Schaffer collaterals (1 Hz, 900 pulses, 10-20 μ A, in D-AP5). The amplitude of mGluR-LTD was significantly greater when high resistance pipettes were employed ($n = 8$ and 5 for low and high resistance pipettes, respectively, $p < 0.001$). 500 μ M (S)-MCPG prevented mGluR-LTD ($n = 5$). (D) Summary of results from extracellular stimulation experiments. *** $p < 0.001$.

baseline, $n = 5$, $p < 0.001$, high versus low resistance pipettes: $p < 0.001$; Fig. 3.1D), indicating that cytoplasmic dialysis significantly disrupts metabotropic signaling. Application of (S)-MCPG (500 μ M) prevented mGluR-LTD (low resistance pipettes: $100 \pm 4\%$ of baseline, $n = 5$, $p = 0.95$; high resistance

Paired recordings
in D-AP5




Figure 3.2 Representative traces obtained during paired recordings for the experimental data plotted in Figure 3.1A. Traces represent averages of 10 sweeps.

pipettes: $103 \pm 6\%$, $n = 5$, $p = 0.6$; Fig. 3.1C). In the relatively young slice cultures used here (3-4 weeks), the PP-LFS protocol did not further enhance mGluR-LTD ($n = 5$, $p = 0.98$), and blocking glutamate transporters with TBOA had no significant effect ($n = 5$, $p = 0.4$; Fig. 3.3).

When Schaffer collaterals are stimulated extracellularly, it can be difficult to rule out that the perforant pathway, other CA1 pyramidal cells, or subicular cells are inadvertently activated. In our slice cultures, however, the subiculum is removed, and perforant path fibers have completely disappeared after two weeks in culture (Zimmer and Gähwiler, 1984). Furthermore, monosynaptic connectivity between CA1 pyramidal cells is 0.8% in acute slices (Thomson & Radpour, 1991), and marginally higher in slice cultures. Nevertheless, we performed experiments in which Schaffer collaterals were activated with the stimulating electrode positioned in the stratum oriens instead of the stratum radiatum of CA3 ($n = 4$), making it unlikely that CA1 pyramidal cell axons are activated. The data show that mGluR-LTD was not significantly different ($p = 0.9$; stratum oriens: $43 \pm 4\%$ of baseline, $n = 4$; stratum radiatum: $42 \pm 3\%$ of baseline, $n = 5$; Fig. 3.4).




Figure 3.3 A comparison of mGluR-LTD induced in CA1 pyramidal cells recorded with high resistance pipettes by extracellular stimulation of Schaffer collaterals using single pulse LFS, PP-LFS, and PP-LFS in the presence of 10 μ M TBOA ($n = 5$ for each protocol). No significant difference in the amplitudes of LTD were observed (LFS: $42 \pm 3\%$ of baseline, $p < 0.001$; PP-LFS: $42 \pm 3\%$ of baseline, $p < 0.001$; PP-LFS + TBOA: $35 \pm 8\%$ of baseline, $p < 0.001$; LFS versus PP-LFS: $p = 0.98$; PP-LFS versus PP-LFS + TBOA: $p = 0.4$).




Figure 3.4 Evidence that a stimulation electrode positioned in CA3 stratum radiatum activates Schaffer collaterals but not other excitatory inputs to CA1. The magnitude of LTD induced by activating Schaffer collaterals with a stimulating placed in CA3 stratum oriens (where CA1 pyramidal cell axons are unlikely to be activated) is not significantly different than when the stimulating electrode is placed in stratum radiatum ($p = 0.9$; $n = 4$ and 5 for stratum oriens and stratum radiatum, respectively).

3.4.2 A single CA3 pyramidal cell does not evoke a somatic current mediated by mGluRs in a connected CA1 pyramidal cell

To examine why mGluR-LTD could not be induced between pyramidal cell pairs, we checked whether an mGluR-mediated current could be evoked in a CA1 pyramidal cell in response to firing of a single connected CA3 pyramidal cell. Following blockade of NMDA, AMPA/kainate, GABA_A and GABA_B receptors, even a high frequency train of 30 APs at 100 Hz failed to evoke a metabotropic response in a CA1 pyramidal cell recorded with a high resistance pipette at -50 mV, a membrane potential where mGluR-mediated currents are maximal (Gee et al., 2003) (n = 3; Fig. 3.5A, lower). Blocking glutamate transporters (TBOA 50 μ M) also did not reveal a metabotropic response (n = 3; Fig. 3.5A, lower). A similar negative result was obtained with the perforated patch-clamp method (n = 3; Fig. 3.6A).

With extracellular stimulation of Schaffer collaterals (100 Hz, 4 pulses, 20-50 μ A, 100 μ s duration), mGluR-mediated EPSCs were consistently evoked in CA1 pyramidal cells, as previously reported (Congar et al., 1997; Nakamura et al., 1999). A metabotropic response was, however, only observed with high resistance recording pipettes (n = 10 and 28 for low and high resistance pipettes, respectively; Fig. 3.5B, upper). The time to peak current (933.3 ± 67.6 ms, n = 28) and the 90-10% decay time (2927.2 ± 187.0 ms, n = 28) of these EPSCs were two orders of magnitude greater than those of ionotropic AMPA EPSCs. Consistent with responses mediated by group I mGluRs, the EPSCs were blocked by (S)-MCPG (500 μ M, n = 13; Fig. 3.5B, lower). Furthermore, TBOA (10 μ M) significantly potentiated mGluR EPSCs evoked by extracellular stimulation ($192 \pm 24\%$ of baseline, n = 6, p < 0.05; Fig. 3.7). Similar results were obtained with the perforated patch-clamp method, with no significant difference in response amplitude obtained with high resistance pipettes (n = 8) versus perforated patch-clamp (n = 5, p > 0.3; Fig. 2C; Fig. 3.6B).




Figure 3.5 Activation of a single CA3 pyramidal cell does not evoke a discernible somatic current mediated by mGluRs in a synaptically connected CA1 pyramidal cell. (A) To enhance mGluR-mediated currents, postsynaptic CA1 cells were voltage-clamped at -50 mV using recording pipettes with a resistance of 10-15 M Ω . Upper, a single AP in a CA3 pyramidal cell elicited a fast EPSC in a CA1 pyramidal cell ($V_h = -70$ mV). Lower, blocking NMDA (50 μ M D-AP5), AMPA/kainate (50 μ M CNQX), GABA_A (100 μ M picrotoxin), and GABA_B receptors (5 μ M CGP 62349) failed to reveal a slow metabotropic current in response to a 100 Hz train of 30 APs. Inhibition of glutamate transporters (50 μ M TBOA) under these conditions also did not reveal an mGluR-mediated current. Traces represent averages of 10 sweeps. (B) Upper, extracellular stimulation of Schaffer collaterals (100 Hz, 4 pulses, 30 μ A, $V_h = -50$ mV) did not evoke an mGluR-mediated response in CA1 pyramidal cells when recording pipettes of 5 M Ω were used. The same stimulation evoked a slow EPSC when using high resistance recording pipettes (12 M Ω). Lower, the slow EPSC was inhibited by (S)-MCPG (500 μ M, $n = 13$). Responses were elicited at 0.1 Hz. Traces represent averages of 5 sweeps. (C) Average amplitudes of mGluR EPSCs evoked with extracellular stimulation ($n = 8$ and 5 for high resistance pipettes and perforated patch-clamp, respectively).




Figure 3.6 Perforated patch-clamp data showing that activation of a single CA3 pyramidal cell does not evoke a current mediated by mGluRs in a synaptically connected CA1 pyramidal cell. (A) Triggering one AP in a CA3 pyramidal cell by current injection elicited an EPSC in a CA1 pyramidal cell ($V_h = -70$ mV). (B) In the presence of CNQX (50 μ M), D-AP5 (50 μ M), picrotoxin (100 μ M), and CGP 62349 (5 μ M), triggering 30 APs at 100 Hz in a CA3 pyramidal cell elicited no current in the connected CA1 pyramidal cell ($V_h = -50$ mV), even when glutamate transporters were blocked with 50 μ M TBOA. Traces represent averages of 10 sweeps. (C) Extracellular stimulation of the Schaffer collaterals evoked a slow EPSC mediated by mGluRs in a CA1 pyramidal cell. The mGluR EPSC amplitude was inhibited by (S)-MCPG (500 μ M, $n = 4$). Responses were elicited at 0.1 Hz. Traces represent averages of 5 sweeps.




Figure 3.7 Application of TBOA potentiated mGluR EPSCs evoked by extracellular stimulation ($10 \mu\text{M}$; $n = 6$). Recording pipettes with resistance of $10\text{-}15 \text{ M}\Omega$ were employed. $*p < 0.05$. Traces represent averages of 5 sweeps.

3.4.3 Estimating the number of Schaffer collaterals required to induce mGluR-mediated responses

As stimulation of a single CA3 pyramidal cell did not induce a metabotropic current in a CA1 pyramidal cell, we performed experiments to estimate the minimum number of synaptic inputs to a CA1 pyramidal cell that must be activated to evoke an mGluR-mediated response. We determined that a threshold stimulation intensity of $20 \mu\text{A}$ applied to the Schaffer collaterals (100 Hz ,






Figure 3.8 Estimating the number of CA3 axons targeting one CA1 pyramidal cell that must be activated to induce an mGluR-mediated current or mGluR-LTD. (A) Average amplitudes of mGluR EPSCs evoked with 2 pulses at 100 Hz were plotted against stimulation intensity (10-50 μ A, V_h = -50 mV, n = 5, high resistance pipettes). (B) AMPA EPSCs in CA1 pyramidal cells elicited with extracellular stimulation of Schaffer collaterals with 1 or 2 pulses at 100 Hz (V_h = -70 mV). Average amplitudes of AMPA EPSCs were plotted against stimulation intensity (n = 7). (C) Paired recordings between CA3-CA1 pyramidal cells. Four APs (100 Hz) triggered in a CA3 pyramidal cell elicited AMPA EPSCs in a connected CA1 pyramidal cell voltage-clamped at -70 mV. Amplitudes of the first EPSCs and the average amplitudes of the first and second EPSCs are shown (n = 22). Outliers may reflect multiquantal release (Tong and Jahr, 1994). (D) mGluR-LTD can be induced in a CA1 pyramidal cell with extracellular stimulation of Schaffer collaterals (1 Hz, 900 pulses, in 50 μ M D-AP5) with a stimulation intensity of 10 μ A, but not 5 μ A (n = 3). The amplitude of the AMPA response, evoked with a stimulation intensity corresponding to the threshold to induce mGluR-LTD or an mGluR-mediated EPSC, was divided by the amplitude of a unitary AMPA response to estimate the number of inputs to a CA1 pyramidal cell required for an mGluR-mediated response.

2 pulses, 100 μ s duration) is necessary to evoke a discernible mGluR-mediated current of ~ 10 pA in CA1 pyramidal cells voltage-clamped at -50 mV ($n = 5$; Fig. 3.8A; Fig. 3.9A). The identical 2-pulse stimulation protocol evoked two AMPA EPSCs, whose mean amplitude was ~ 470 pA at 20 μ A, derived from the exponential fit of the curve ($f = y_0 + a \cdot (1 - \exp(-b \cdot x))$); Fig. 3.8B; Fig. 3.9B). As the median amplitude of the two unitary AMPA responses was 23.1 pA ($n = 22$ pairs; Fig. 3.8C; Fig. 3.9C), we can estimate that ~ 20 CA3 synaptic inputs released glutamate to yield a response of 470 pA. Therefore, approximately 20 CA3 pyramidal cell axons targeting a given CA1 pyramidal cell must be activated concurrently to induce a somatic current mediated by group I mGluRs. We then checked whether the threshold stimulation intensity for mGluR-LTD is similar to that required to induce an mGluR-mediated current. We found that a much lower stimulation intensity of 10 μ A (at 1 Hz) was sufficient to induce mGluR-LTD. At a still lower stimulation intensity of 5 μ A, an AMPA receptor-mediated current was no longer observed. As glutamate binds with an approximately 30-fold higher affinity to mGluRs (Conn and Pin, 1997) than to AMPA receptors (Jonas and Sakmann, 1992), it is nonetheless possible that this lower stimulation intensity might induce mGluR-LTD. However, when stimulation intensity was reduced to 5 μ A, LTD was no longer observed (5 μ A: $101 \pm 4\%$ of baseline, $p = 0.9$; 10 μ A: $46 \pm 7\%$ of baseline, $p < 0.05$; $n = 3$; Fig. 3.8D). Thus, an intensity of 10 μ A is close to threshold for inducing mGluR-LTD. A stimulus of 10 μ A evokes an extracellular AMPA receptor-mediated response of ~ 190 pA (Fig. 3.8B), which when divided by the median value of a single unitary AMPA response (23.3 pA, $n = 22$ pairs; Fig. 3.8C) provides an estimate of 8 active CA3 inputs required to induce mGluR-LTD in a targeted CA1 pyramidal cell.

A mGluR EPSC Extracellular stimulation



B AMPA EPSC Extracellular stimulation



C AMPA EPSC Paired recordings




Figure 3.9 Representative traces obtained during recordings for the experimental data plotted in Fig. 3.8. (A) an mGluR EPSC evoked by extracellular stimulation of Schaffer collaterals (100 Hz, 2 pulses, 20 μ A, V_h = -50 mV). (B) AMPA EPSC evoked by extracellular stimulation of Schaffer collaterals (100 Hz, 2 pulses, 20 μ A, V_h = -70 mV). (C) AMPA EPSCs in paired recordings between CA3-CA1 pyramidal cells. Traces from two different pairs are shown. Four APs (100 Hz) in a CA3 pyramidal cell elicited unitary AMPA EPSCs in a connected CA1 pyramidal cell voltage-clamped at -70 mV. Traces represent averages of 5, 5 and 20 sweeps in (A), (B) and (C), respectively.

3.5 Discussion

The main finding of this study is that in contrast to NMDAR-LTD (Debanne et al., 1996), mGluR-LTD assessed by somatic recording in CA1 pyramidal cells could not be induced by stimulating a single synaptically connected CA3 pyramidal cell. A number of possibilities may account for this difference. First, NMDA receptors in CA1 pyramidal cells are usually located in the middle of the synaptic disk (Racca et al., 2000), whereas metabotropic receptors display a perisynaptic or extrasynaptic localization (Lujan et al., 1996). Thus, cooperative presynaptic activity could be necessary to allow pooling and diffusion of neurotransmitter to the more distantly positioned metabotropic receptors, as demonstrated for activation of GABA_B receptors (Scanziani, 2000). This mechanism, however, does not appear to be involved in inducing mGluR current, as inhibition of glutamate transporters with TBOA to promote synaptic spillover of neurotransmitter did not reveal a response. A second possibility is that as opposed to NMDA receptors, which are expressed in virtually all spines in CA1 pyramidal cells, (Takumi et al., 1999; Racca et al., 2000), the expression of dendritic mGluRs may be relatively low, such that diffusion of transmitter from a single release site will rarely reach an mGluR-positive spine. Evidence for a sparse distribution of functional dendritic mGluRs is provided in a recent study showing that metabotropic responses in CA1 pyramidal cells could only be induced in a subset of endoplasmic reticulum-containing mushroom spines (Holbro et al., 2009), which represent less than 10% of all spines. Furthermore, the failure rate with glutamate uncaging for mGluR-mediated responses in mGluR-positive spines was approximately 80%. Even though a single CA3 pyramidal cell can make multiple functional contacts with a synaptically coupled CA1 pyramidal cell (Hsia et al., 1998) – estimated at between 2 to 4 (Sorra and Harris, 1993) – the probability that one of these few contacts will produce a metabotropic response is therefore very low. Indeed, our experiments indicate that ~20 CA3 pyramidal cell inputs to a CA1 pyramidal cell must be activated for reliable induction of somatic mGluR-mediated current. This is, nevertheless, a

relatively small number considering that a CA1 pyramidal cell *in vivo* receives, on average, input from 5,500 CA3 pyramidal cells (Amaral et al., 1990). Moreover, we estimate that even fewer CA3 inputs (less than ten) must be activated to induce mGluR-LTD. When an mGluR-positive spine is first identified and then specifically stimulated by two-photon glutamate uncaging, it is, however, possible to obtain mGluR-LTD (Holbro et al., 2009). This difference to our experiments is likely to reflect the fact that prior identification of mGluR-positive synapses is not possible at present with paired recording, and, as there are multiple functional contacts between CA3-CA1 pyramidal cell pairs, LTD of at best one of these synapses may not be discernible within the compound synaptic response. Furthermore, localized uncaging of glutamate may be more efficient than synaptically released glutamate at reaching peripherally localized mGluRs.

A lower threshold for mGluR-LTD as compared to mGluR-mediated current in CA1 pyramidal cells could reflect differences in the respective intracellular transduction mechanisms. Thus, an mGluR-dependent calcium signal to initiate LTD induction may have a lower stimulation threshold than an mGluR-mediated current, as demonstrated in cerebellar Purkinje cells (Finch and Augustine, 1998; Takechi et al., 1998). It has not been possible to repeat this type of experiment in CA3-CA1 paired recordings because of the difficulty in localizing the very few activated spines in the large dendritic tree of a pyramidal cell.

A second finding emerging from our study is the pronounced sensitivity of mGluR-mediated responses to cytoplasmic dialysis. This phenomenon may explain why in most studies using whole-cell patch recording, concentrations of agonists to activate postsynaptic mGluRs exceed EC_{50} values for group I mGluRs (Conn and Pin, 1997) by one to two orders of magnitude. The rundown of metabotropic responses can be avoided with the perforated patch-clamp method (Horn and Marty, 1988), but this approach is technically demanding. Here, we show that a simpler method employing patch pipettes with a higher resistance (10 to 15 M Ω) allows mGluR-mediated currents to be recorded with similar amplitudes to those obtained with the perforated patch-clamp technique.

In conclusion, we have shown that even under conditions where cytoplasmic dialysis is minimized, mGluR-LTD is not observed in CA3-CA1 pyramidal cell pairs. Temporally contiguous input from multiple Schaffer collaterals must target a CA1 pyramidal cell to induce mGluR-mediated currents and mGluR-LTD. Interestingly, the requirement for multiple inputs does not reflect a cooperative mechanism involving synaptic spillover of glutamate (Arnth-Jensen et al., 2002), but rather is consistent with a sparse expression of functional mGluRs in the dendritic tree of pyramidal cells.

Chapter 4

Group I metabotropic glutamate receptors modulate hippocampal rhythmic activity

4.1 Abstract

In this study, we check the activation conditions of group I metabotropic glutamate receptor-mediated currents in hippocampal interneurons. We show that activation of a single CA3 pyramidal cell does not evoke a somatic current mediated by group I mGluRs in a synaptically connected interneuron. Hippocampal interneurons are required for induction and propagation of theta rhythm. Our observations show that group I mGluRs in hippocampal interneurons are activated during theta oscillations and modulate the frequency of these oscillations.

4.2 Introduction

In the behaving animals, synchronous oscillations of the membrane potential of large neuronal populations can occur at various frequencies in the hippocampus. Theta rhythm (4-15 Hz) (Bland 1986; Lopes da Silva et al. 1990; Stewart and Fox 1990; Vinogradova 1995; Vertes and Kocsis 1997; Kamondi et al., 1998; Harris et al., 2001; Buzsáki, 2002) is present mainly during locomotion and other voluntary movements (Grastyán et al., 1959; Vanderwolf, 1969) and rapid eye movement (REM) sleep (Jouvet, 1969). Theta waves have been also assumed to carry mnemonic processes (Lisman and Idiart 1995; Raghavachari et al. 2001). Loss of hippocampal theta rhythm results in spatial memory deficits in the rat (Winson 1978).

Mechanisms of the generation of theta rhythm in the hippocampus are not completely understood yet. However, it is clear that the reciprocal connection between the medial septum-diagonal band of Broca (MS-DBB) and the hippocampus plays an essential role (Green and Arduni, 1954; Petsche et al., 1962; Lawson and Bland, 1993; Sik et al., 1994; Markowska et al., 1995; King et

al., 1998; Dragoi et al., 1999; Buzsáki 2002; Gulyas et al., 2003). Hippocampal interneurons represent the only output from the hippocampus to the MS-DBB (Toth et al., 1993) and are required for the induction and propagation of theta rhythm (Buzsáki et al., 1983; Sik et al., 1994; Freund and Antal, 1988; Buzsáki and Chrobak, 1995; Cobb et al., 1995; Dragoi et al., 1999; Buzsáki 2002; Gulyas et al., 2003; Mann and Paulsen, 2007). One hippocampal interneuron innervates more than 1,000 pyramidal cells (Halasy et al., 1996). Thus, they are in a key position to synchronize the activity of pyramidal cells.

Theta rhythm can be mimicked *in vitro* by activation of muscarinic acetylcholine receptors (mAChRs) (Konopacki et al., 1987). In hippocampal slice cultures, rhythmic activity can be induced with very low (10-20 nM) concentrations of the mAChR agonist methacholine (Fische et al., 1999, 2002). In this study, we show that group I mGluRs in hippocampal interneurons are activated during theta oscillations and modulate the frequency of these oscillations.

4.3 Materials and Methods

4.3.1 Slice cultures and electrophysiology

For the preparation of hippocampal organotypic slice cultures, see chapter 2.

Whole-cell voltage-clamp recordings were obtained with an Axopatch 200B amplifier (Molecular Devices). Recording pipettes were filled with (in mM) 120 K-gluconate, 10 L-glutamic acid, 5 KCl, 10 Hepes, 10 EGTA, 2 MgATP, 5 creatine phosphate (CrP), 0.4 NaGTP, 0.07 CaCl₂ (pH 7.2, ~290 mOsm). Membrane potentials were corrected for liquid junction potentials. Series resistance and input resistance were monitored regularly. Currents were filtered at 5 kHz and

analyzed off-line (pClamp 9, Molecular Devices). All numerical data are presented as the mean \pm SEM. Significance was tested using two-tailed t-test.

4.3.2 Extracellular stimulation

For monopolar extracellular stimulation, glass pipettes filled with ACSF were placed 100-200 μ m away from the recorded cell to stimulate Schaffer collaterals. To evoke a current mediated by group I mGluRs, a single-pulse stimulation (10-50 μ A, 100 μ s) was delivered. CA1 interneurons were clamped at -50 mV. AMPA, NMDA, GABA_A and GABA_B receptors were blocked by CNQX (50 μ M), D-AP5 (100 μ M), picrotoxin (100 μ M) and CGP 62349 (5 μ M), respectively.

4.3.3 Paired recordings

For current-clamp recording, pipettes (3-5 M Ω) were filled with (in mM) 120 K-gluconate, 10 L-glutamic acid, 5 KCl, 10 Hepes, 1 EGTA, 2 MgATP, 5 creatine phosphate, 0.4 NaGTP, 0.07 CaCl₂ (pH 7.2, ~290 mOsm). For voltage-clamp recordings, pipettes (10-15 M Ω) were filled with (in mM) 126.6 CsF, 8.4 CsCl, 10 Hepes, 10 EGTA, 2 MgATP, 5 creatine phosphate, 0.4 NaGTP (pH 7.2, ~290 mOsm).

4.3.4 Drugs

CNQX (6-cyano-7-nitroquinoxaline-2,3-dione), D-AP5 (D-(-)-2-amino-5-phosphonopentanoic acid), (S)-MCPG ((S)- α -methyl-4-carboxyphenylglycine) and MPEP (2-methyl-6-(phenylethynyl)pyridine hydrochloride) were purchased from Ascent Scientific (Bristol, UK). DL-TBOA (DL-threo-b-benzyloxyaspartic acid) and YM 29819 was purchased from Tocris (Bristol, UK). CGP 62349 was kindly provided by Novartis (Basel, Switzerland). All other chemicals were purchased from Sigma/Fluka.


4.4 Results

4.4.1 Activation of a single CA3 pyramidal cell does not evoke a somatic current mediated by group I mGluRs in a synaptically connected interneuron

We checked whether a group I mGluR-mediated current could be activated in a hippocampal interneuron in stratum radiatum or stratum oriens of the CA3 region when only a single presynaptically connected CA3 pyramidal cell is evoked to fire. High resistance pipettes (10-15 M Ω) were employed to minimize cytoplasmic dialysis, which has been shown to be critical for induction of group I mGluR-mediated responses (see chapter 3). Following the blockade of NMDA, AMPA/kainate, GABA_A and GABA_B receptors, even a high-frequency train of 30 APs at 100 Hz failed to evoke a metabotropic response in an interneuron voltage-clamped at -50 mV, a potential where mGluR-mediated currents are maximal (Gee et al., 2003) (n = 3; Fig. 4.1, lower). As postsynaptic mGluRs in hippocampal interneurons are localized in perisynaptic and extrasynaptic zones (Lujan et al., 1996, 1997), membrane transporters may prevent glutamate access to mGluRs. However, in the presence of the glutamate transporter inhibitor TBOA (50 μ M) (Jabaudon et al., 1999), a metabotropic response still could not be evoked (n = 3; Fig. 4.1, lower).

Paired recordings

High resistance pipettes



100 Hz, 30 APs





Figure 4.1 Activation of a single CA3 pyramidal cell does not evoke a current mediated by group I mGluRs in a connected interneuron of the CA3 region. Postsynaptic interneurons were voltage-clamped at -50 mV and recording pipettes with a resistance of 10-15 M Ω were used. Upper, one AP triggered in a CA3 pyramidal cell elicited a fast EPSC in a connected CA1 interneuron ($V_h = -70$ mV). Lower, blocking ionotropic receptors (100 μ M D-AP5, 50 μ M CNQX) and GABA_A and GABA_B receptors (100 μ M picrotoxin, 5 μ M CGP 62349) failed to reveal a slow metabotropic current in response to a 100 Hz train of 30 APs. Inhibition of glutamate transporters (50 μ M TBOA) under these conditions also did not reveal an mGluR-mediated current. Traces represent averages of 10 sweeps.

4.4.2 Extracellular stimulation of Schaffer collaterals evokes group I mGluR-mediated EPSCs in interneurons of the CA1 region

With extracellular stimulation of Schaffer collaterals (100 Hz, 4 pulses, 20-50 μ A, 100 μ s duration), mGluR-mediated EPSCs were consistently evoked in interneurons of the CA1 region ($n = 4$; Fig. 4.2A, B). These EPSCs were blocked by the group I mGluR antagonist (S)-MCPG (500 μ M, $n = 3$; Fig. 4.2B).




Figure 4.2 Extracellular stimulation of Schaffer collaterals evokes group I mGluR-mediated EPSCs in interneurons of the CA1 region. (A) Upper, extracellular stimulation of Schaffer collaterals (100 Hz, 4 pulses, 10-50 μ A, $V_h = -50$ mV) evoked an mGluR-mediated current in CA1 interneurons when using high resistance recording pipettes (10-15 M Ω). Lower, the slow EPSC was significantly reduced by (S)-MCPG (500 μ M, $n = 3$). Responses were elicited at 0.1 Hz. Traces represent averages of 5 sweeps. (B) Average amplitudes of mGluR EPSCs evoked with extracellular stimulation (100 Hz, 2 pulses, 10-50 μ A, $V_h = -50$ mV; $n = 4$).

4.4.3 Group I mGluRs modulate hippocampal theta rhythm

More than 1,000 pyramidal cells terminate on a single interneuron in hippocampus (Hayashi et al., 1993). Although a single CA3 pyramidal cell does not activate group I mGluRs in a synaptically connected interneuron, concurrent activation of several pyramidal cells may activate group I mGluRs in hippocampal interneurons. During theta oscillations approximately 2% of pyramidal cells fire together (Csicsvari et al. 1998). Therefore, we hypothesize that group I mGluRs in hippocampal interneurons are activated during theta oscillations.

Rhythmic activity was induced with 50 nM concentration of the muscarinic acetylcholine receptor agonist methacholine (Fische et al., 1999, 2002). Application of mGluR1 antagonist YM 298198 (1 μ M) and mGluR5 antagonist MPEP (10 μ M) decreased the frequency of theta oscillations (for YM 298198, $61 \pm 7\%$ of baseline, $n = 7$, $p < 0.05$; for MPEP, $51 \pm 4\%$ of baseline, $n=9$, $p < 0.001$; Figure 4.3). The above data show that group I mGluRs in hippocampal interneurons are activated during theta oscillations.




Figure 4.3 Group I metabotropic glutamate receptors modulate hippocampal rhythmic activity. (A) Application of the mGluR5 antagonist MPEP (10 μ M) decreased the frequency of oscillations as illustrated in the autocorrelograms. Inset, representative traces recorded from a CA1 pyramidal cell voltage clamped at -70 mV. (b) The frequency of the oscillations was decreased by the mGluR1 antagonist YM 298198 (1 μ M, $n = 7$) and by the mGluR5 antagonist MPEP (10 μ M, $n = 9$); * $p < 0.05$ and *** $p < 0.001$.

4.5 Discussion

In this study, we report that activation of a single CA3 pyramidal cell does not evoke a somatic current mediated by group I mGluRs in a synaptically connected interneuron. However, mGluR-mediated EPSCs were consistently evoked with extracellular stimulation of Schaffer collaterals. This is reminiscent of the result in chapter 3, which is, concurrent activation of ~8 CA3 pyramidal cells is required to activate group I mGluRs in a CA1 pyramidal. Therefore, we checked whether concurrent activation of several pyramidal cells are necessary for activation of group I mGluRs in interneurons. We show that block of group I mGluRs modulates the frequency of theta oscillations, which proves that group I mGluRs in interneurons are activated by synchronized activity of pyramidal cells during hippocampal theta rhythm.

Although mGluR1 and mGluR5 are highly homologous, their function (Valenti et al., 2002) and distribution (see chapter 1.3) are reported to be different. Our data show that blockade of mGluR1 or mGluR5 can reduce the frequency of hippocampal theta oscillations.

Hippocampal interneurons are required for the induction and propagation of theta rhythm (Buzsáki et al., 1983; Sik et al., 1994; Freund and Antal, 1988; Buzsáki and Chrobak, 1995; Cobb et al., 1995; Dragoi et al., 1999; Buzsáki 2002; Gulyas et al., 2003; Mann and Paulsen, 2007). However, rhythmic activity relies on the balance between inhibition and excitation (Shu et al., 2003; Mann and Paulsen, 2007). It would be interesting to compare the functions of group I mGluRs located in pyramidal cells and in interneurons. Up to now, there is no pharmacological tool that can selectively block or activate group I mGluRs selectively only in pyramidal cells or only in interneurons. Transgenic mice in which the mGluR1 or/and mGluR5 coding gene is knocked out selectively from pyramidal cells or from interneurons would be useful tools to study this question.

Chapter 5

Conclusion and perspectives

5.1 Conclusion and discussion

My thesis project was to determine the conditions required for activation of group I mGluRs in rat hippocampus.

Firstly, as described in chapter 2, I checked in hippocampal slice cultures whether the ambient glutamate concentration in the extracellular space (typically 2 μ M in the hippocampal slices) is sufficient to tonically activate group I mGluRs (whose EC_{50} is ~ 10 μ M). However, I show that group I mGluRs in CA1 pyramidal cells are not tonically activated by ambient glutamate in hippocampal slice cultures. One possible explanation is that the background level of extracellular glutamate in hippocampal organotypic slice cultures is lower than in the brain *in vivo* or in acute brain slices. Another possible explanation is that the ambient concentration of glutamate in acute hippocampal slices is much lower than the value formerly reported. In a new measurement, the ambient concentration of glutamate in acute hippocampal slices was reported to be ~ 25 nM (Herman and Jahr, 2007).

In chapter 3, I present my investigation on the electrical stimulation conditions to activate group I mGluRs in CA1 pyramidal cells. NMDAR-LTD and mGluR-LTD have both been found in CA1 pyramidal cells. For the induction of NMDAR-LTD, cooperative interactions of multiple presynaptic inputs are not essential, such that appropriately timed discharge of a single CA3 pyramidal cell can lead to homosynaptic LTD in a targeted CA1 pyramidal cell (Debanne et al., 1996). Activation of group I mGluRs with low frequency stimulation of many Schaffer collaterals can induce mGluR-LTD in CA1 pyramidal cells. In chapter 3, I show that activation of a single CA3 pyramidal cell does not induce mGluR-LTD in a connected CA1 pyramidal cell. An estimate based on data obtained from paired recording and extracellular stimulation indicates that 8 active CA3 inputs are required to induce mGluR-LTD in a targeted CA1 pyramidal cell. Similarly, I show that activation of a single CA3 pyramidal cell does not evoke a somatic current

mediated by mGluRs in a connected CA1 pyramidal cell. An estimate based on data obtained from paired recording and extracellular stimulation indicates that 20 CA3 pyramidal cell axons targeting a given CA1 pyramidal cell must be activated concurrently to induce a somatic current mediated by group I mGluRs. This may be due to the sparse distribution of functional dendritic group I mGluRs, which was shown in a recent study by Holbro et al. (2009). The authors showed that responses mediated by group I mGluR in CA1 pyramidal cells could only be induced in a subset of endoplasmic reticulum-containing mushroom spines, which represent less than 10% of all spines. Furthermore, the failure rate with glutamate uncaging for mGluR-mediated responses in these mGluR-positive spines was approximately 80%.

I show that 8 and 20 CA3 inputs are required to induce mGluR-LTD and an mGluR-mediated current in a targeted CA1 pyramidal cell, respectively. This is, actually, a relatively small number considering that a CA1 pyramidal cell *in vivo* receives, on average, input from 5,500 CA3 pyramidal cells (Amaral et al., 1990).

In my investigation of the conditions to activate mGluR-mediated responses in CA1 pyramidal cells, an interesting finding was the pronounced sensitivity of mGluR-mediated responses to cytoplasmic dialysis associated with conventional whole-cell patch clamp recording. The rundown of metabotropic responses can be avoided by using the perforated patch-clamp method (Horn and Marty, 1988), but this approach is technically very demanding. Here, I show that a simpler method, in which patch pipettes with a resistance of 10 to 15 M Ω instead of 3 to 5 M Ω are employed, allow mGluR-mediated currents to be recorded with similar amplitudes to those obtained with the perforated patch-clamp technique.

Finally, in chapter 4, I check the activation conditions of group I metabotropic glutamate receptor-mediated currents in hippocampal interneurons. I show that although extracellular stimulation of Schaffer collaterals evokes group I mGluR-mediated EPSCs in interneurons of the CA1 region, activation of a single CA3

pyramidal cell does not evoke a somatic current mediated by group I mGluRs in a synaptically connected interneuron. This is reminiscent of the observations in chapter 3, in which I found that concurrent activation of CA3 pyramidal cells is required to activate group I mGluRs in a CA1 pyramidal cell. I find that block of group I mGluRs modulates the frequency of hippocampal theta oscillations, which proves that group I mGluRs in interneurons are activated by synchronized activity of pyramidal cells during hippocampal theta rhythm. This finding is consistent with the fact that more than 1,000 pyramidal cells terminate on a single interneuron in hippocampus (Hayashi et al., 1993) and during theta oscillations approximately 2% of pyramidal cells fire together (Csicsvari et al. 1998).

5.2 Perspectives

In the study of Holbro et al., 2009, taking advantage of a two-photon glutamate uncaging method, the authors can identify mGluR-positive spines and then specifically stimulate an identified spine to obtain mGluR-LTD. However, it has not been possible to identify mGluR-positive synapses with paired recordings. Furthermore, localized glutamate uncaging may be more efficient than synaptically released glutamate at reaching peripherally localized mGluRs. In chapter 3, I show that less CA3 inputs (8 v.s. 20) are required to induce mGluR-LTD than a somatic mGluR-mediated current in a targeted CA1 pyramidal cell. An mGluR-dependent calcium signal to initiate LTD induction may have a lower stimulation threshold than an mGluR-mediated current, as has been demonstrated in cerebellar Purkinje cells (Finch and Augustine, 1998; Takechi et al., 1998). At present, it is not possible to repeat this type of experiment in CA3-CA1 paired recordings as it is difficult to localize the very few activated spines in the large dendritic tree of a CA1 pyramidal cell. Two-photon glutamate uncaging can be a practical method to address this question.

Hippocampal interneurons are required for the induction and propagation of theta rhythm (Buzsáki et al., 1983; Sik et al., 1994; Freund and Antal, 1988; Buzsáki and Chrobak, 1995; Cobb et al., 1995; Dragoi et al., 1999; Buzsáki 2002; Gulyas et al., 2003; Mann and Paulsen, 2007). However, rhythmic activity relies on the balance between inhibition and excitation (Shu et al., 2003; Mann and Paulsen, 2007). It would be interesting to compare the functions of group I mGluRs located in pyramidal cells and in interneurons. Up to now, there is no pharmacological tool that can selectively block or activate group I mGluRs selectively only in pyramidal cells or only in interneurons. Transgenic mice in which the mGluR1 or/and mGluR5 coding gene is knocked out selectively from pyramidal cells or from interneurons would be useful tools to study this question.

References

- Abdul-Ghani MA, Valiante TA, Carlen PL, Pennefather PS (1996) Metabotropic glutamate receptors coupled to IP₃ production mediate inhibition of IAHP in rat dentate granule neurons. *J Neurophysiol* 76:2691-2700.
- Abe T, Sugihara H, Nawa H, Shigemoto R, Mizuno N (1992) Molecular characterization of a novel metabotropic glutamate receptor mGluR5 coupled to inositol phosphate/Ca²⁺ signal transduction. *J Biol Chem* 267:13361.
- Acuna-Goycolea C, Li Y, van den Pol AN (2004) Group III metabotropic glutamate receptors maintain tonic inhibition of excitatory synaptic input to hypocretin/orexin neurons. *J Neurosci* 24:3013-3022.
- Ahmed MS, Siegelbaum SA (2009) Recruitment of N-Type Ca²⁺ Channels during LTP Enhances Low Release Efficacy of Hippocampal CA1 Perforant Path Synapses. *Neuron* 63:372-385.
- Aiba A, Chen C, Herrup K, Rosenmund C, Stevens CF, Tonegawa S (1994) Reduced hippocampal long-term potentiation and context-specific deficit in associative learning in mGluR1 mutant mice. *Cell* 79:365-375.
- Amaral DG, Ishizuka N, Claiborne B (1990) Neurons, numbers and the hippocampal network. *Prog Brain Res* 83:1-11.
- Angulo MC, Kozlov AS, Charpak S, Audinat E (2004) Glutamate released from glial cells synchronizes neuronal activity in the hippocampus. *J Neurosci* 24:6920-6927.
- Ango F, Pin J-P, Tu JC, Xiao B, Worley PF, Bockaert J, Fagni L (2000) Dendritic and axonal targeting of type 5 metabotropic glutamate receptor is regulated by homer1 proteins and neuronal excitation. *J Neurosci* 20:8710-8716.
- Aniksztejn L, Bregestovski P, Ben-Ari Y (1991) Selective activation of quisqualate metabotropic receptor potentiates NMDA but not AMPA responses. *Eur J Pharmacol* 205:327.

- Aniksztejn L, Otani S, Ben-Ari Y (1992) Quisqualate Metabotropic Receptors Modulate NMDA Currents and Facilitate Induction of Long-Term Potentiation Through Protein Kinase C. *European Journal of Neuroscience* 4:500-505.
- Anwyl R (1989) Protein kinase C and long-term potentiation in the hippocampus. *Trends in Pharmacological Sciences* 10:236-239.
- Anwyl R (1999) Metabotropic glutamate receptors: electrophysiological properties and role in plasticity. *Brain Research Reviews* 29:83-120.
- Anwyl R (2006) Induction and expression mechanisms of postsynaptic NMDA receptor-independent homosynaptic long-term depression. *Progress in Neurobiology* 78:17-37.
- Anwyl R (2009) Metabotropic glutamate receptor-dependent long-term potentiation. *Neuropharmacology* 56:735-740.
- Arnth-Jensen N, Jabsaudon D, Scanziani M (2002) Cooperation between independent hippocampal synapses is controlled by glutamate uptake. *Nat Neurosci* 5:325-331.
- Attwell D, Gibb A (2005) Neuroenergetics and the kinetic design of excitatory synapses. *Nat Rev Neurosci* 6:841-849.
- Awad H, Hubert GW, Smith Y, Levey AI, Conn PJ (2000) Activation of metabotropic glutamate receptor 5 has direct excitatory effects and potentiates NMDA receptor currents in neurons of the subthalamic nucleus. *J Neurosci* 20:7871-7879.
- Bashir ZI, Collingridge GL (1992) Synaptic plasticity: long-term potentiation in the hippocampus. *Current Opinion in Neurobiology* 2:328-335.
- Bashir ZI, Bortolotto ZA, Davies CH, Berretta N, Irving AJ, Seal AJ, Henley JM, Jane DE, Watkins JC, Collingridge GL (1993a) Induction of LTP in the hippocampus needs synaptic activation of glutamate metabotropic receptors. *Nature* 363:347-350.
- Bashir ZI, Jane DE, Sunter DC, Watkins JC, Collingridge GL (1993b) Metabotropic glutamate receptors contribute to the induction of long-term

- depression in the CA1 region of the hippocampus. *European Journal of Pharmacology* 239:265-266.
- Baskys A, Malenka RC (1991) Agonists at metabotropic glutamate receptors presynaptically inhibit EPSCs in neonatal rat hippocampus. *The Journal of Physiology* 444:687-701.
- Baude A, Nusser Z, Roberts JDB, Mulvihill E, Jeffrey McIlhinney RA, Somogyi P (1993) The metabotropic glutamate receptor (mGluRI α) is concentrated at perisynaptic membrane of neuronal subpopulations as detected by immunogold reaction. *Neuron* 11:771-787.
- Bayazitov IT, Richardson RJ, Fricke RG, Zakharenko SS (2007) Slow Presynaptic and Fast Postsynaptic Components of Compound Long-Term Potentiation. *J Neurosci* 27:11510-11521.
- Bear MF, Abraham WC (1996) Long-Term Depression in Hippocampus. *Annual Review of Neuroscience* 19:437-462.
- Bear MF, Malenka RC (1994) Synaptic plasticity: LTP and LTD. *Current Opinion in Neurobiology* 4:389-399.
- Bear MF, Huber KM, Warren ST (2004) The mGluR theory of fragile X mental retardation. *Trends in Neurosciences* 27:370-377.
- Benquet P, Gee CE, Gerber U (2002) Two distinct signaling pathways upregulate NMDA receptor responses via two distinct metabotropic glutamate receptor subtypes. *J Neurosci* 22:9679-9686.
- Benveniste H, Drejer J, Schousboe A, Diemer NH (1984) Elevation of the extracellular concentrations of glutamate and aspartate in rat hippocampus during transient cerebral ischemia monitored by intracerebral microdialysis. *Journal of Neurochemistry* 43:1369-1374.
- Bergles DE, Diamond JS, Jahr CE (1999) Clearance of glutamate inside the synapse and beyond. *Current Opinion in Neurobiology* 9:293-298.
- Berthele A, Laurie DJ, Platzer S, Zieglgänsberger W, Tölle TR, Sommer B (1998) Differential expression of rat and human type 1 metabotropic glutamate receptor splice variant messenger RNAs. *Neuroscience* 85:733-749.

- Berthele A, Boxall SJ, Urban A, Anneser JMH, Zieglgänsberger W, Urban L, Tölle TR (1999) Distribution and developmental changes in metabotropic glutamate receptor messenger RNA expression in the rat lumbar spinal cord. *Developmental Brain Research* 112:39-53.
- Bettler B, Mulle C (1995) AMPA and kainate receptors. *Neuropharmacology* 34:123-139.
- Bhave G, Karim F, Carlton SM, Gereau RW (2001) Peripheral group I metabotropic glutamate receptors modulate nociception in mice. *Nat Neurosci* 4:417-423.
- Bianchi L, Della Corte L, Tipton KF (1999) Simultaneous determination of basal and evoked output levels of aspartate, glutamate, taurine and 4-aminobutyric acid during microdialysis and from superfused brain slices. *Journal of Chromatography B: Biomedical Sciences and Applications* 723:47-59.
- Bland BH (1986) The physiology and pharmacology of hippocampal formation theta rhythms. *Progress in Neurobiology* 26:1-54.
- Bleakman D, Rusin KI, Chard PS, Glaum SR, Miller RJ (1992) Metabotropic glutamate receptors potentiate ionotropic glutamate responses in the rat dorsal horn. *Mol Pharmacol* 42:192-196.
- Bleakman D (1999) Kainate receptor pharmacology and physiology. *Cellular and Molecular Life Sciences* 56:558-566.
- Bliss TV, Collingridge GL (1993) A synaptic model of memory: long-term potentiation in the hippocampus. *Nature* 361:31 - 39.
- Bolshakov VY, Siegelbaum SA (1994) Postsynaptic induction and presynaptic expression of hippocampal long-term depression. *Science* 264:1148-1152.
- Bordi F, Ugolini A (1999) Group I metabotropic glutamate receptors: implications for brain diseases. *Progress in Neurobiology* 59:55-79.
- Bortolotto ZA, Collingridge GL (1993) Characterisation of LTP induced by the activation of glutamate metabotropic receptors in area CA1 of the hippocampus. *Neuropharmacology* 32:1-9.

- Bortolotto ZA, Fitzjohn SM, Collingridge GL (1999) Roles of metabotropic glutamate receptors in LTP and LTD in. *Current Opinion in Neurobiology* 9:299-304.
- Brager DH, Johnston D (2007) Plasticity of intrinsic excitability during long-term depression is mediated through mGluR-dependent changes in I(h) in hippocampal CA1 pyramidal neurons. *J Neurosci* 27:13926-13937.
- Bragin A, Jando G, Nadasdy Z, Hetke J, Wise K, Buzsaki G (1995) Gamma (40-100 Hz) oscillation in the hippocampus of the behaving rat. *J Neurosci* 15:47-60.
- Brakeman PR, Lanahan AA, O'Brien R, Roche K, Barnes CA, Huganir RL, Worley PF (1997) Homer: a protein that selectively binds metabotropic glutamate receptors. *Nature* 386:284-288.
- Bredt DS, Nicoll RA (2003) AMPA Receptor Trafficking at Excitatory Synapses. *Neuron* 40:361-379.
- Bruno V, Battaglia G, Copani A, D'Onofrio M, Di Iorio P, De Blasi A, Melchiorri D, Flor PJ, Nicoletti F (2001) Metabotropic glutamate receptor subtypes as targets for neuroprotective drugs. *J Cereb Blood Flow Metab* 21:1013-1033.
- Bruno V, Battaglia G, Ksiazek I, van der Putten H, Catania MV, Giuffrida R, Lukic S, Leonhardt T, Inderbitzin W, Gasparini F, Kuhn R, Hampson DR, Nicoletti F, Flor PJ (2000) Selective activation of mGlu4 metabotropic glutamate receptors is protective against excitotoxic neuronal death. *J Neurosci* 20:6413-6420.
- Buzsáki G, Lai-Wo S L, Vanderwolf CH (1983) Cellular bases of hippocampal EEG in the behaving rat. *Brain Research Reviews* 6:139-171.
- Buzsáki G, Chrobak JJ (1995) Temporal structure in spatially organized neuronal ensembles: a role for interneuronal networks. *Current Opinion in Neurobiology* 5:504-510.
- Buzsáki G (2002) Theta Oscillations in the Hippocampus. *Neuron* 33:325-340.
- Calabresi P, Centonze D, Pisani A, Bernardi G (1999) Metabotropic Glutamate Receptors and Cell-Type-Specific Vulnerability in the Striatum: Implication

- for Ischemia and Huntington's Disease. *Experimental Neurology* 158:97-108.
- Catania MV, Aronica E, Sortino MA, Canonico PL, Nicoletti F (1991) Desensitization of metabotropic glutamate receptors in neuronal cultures. *Journal of Neurochemistry* 56:1329-1335.
- Catania MV, Landwehrmeyer GB, Testa CM, Standaert DG, Penney JJB, Young AB (1994) Metabotropic glutamate receptors are differentially regulated during development. *Neuroscience* 61:481-495.
- Catania MV, Bellomo M, Di Giorgi-Gerevini V, Seminara G, Giuffrida R, Romeo R, De Blasi A, Nicoletti F (2001) Endogenous activation of group-I metabotropic glutamate receptors is required for differentiation and survival of cerebellar purkinje cells. *J Neurosci* 21:7664-7673.
- Cavelier P, Attwell D (2005) Tonic release of glutamate by a DIDS-sensitive mechanism in rat hippocampal slices. *J Physiol* 564:397-410.
- Cavelier P, Hamann M, Rossi D, Mobbs P, Attwell D (2005) Tonic excitation and inhibition of neurons: ambient transmitter sources and computational consequences. *Progress in Biophysics and Molecular Biology* 87:3-16.
- Chapman AG, Nanan K, Williams M, Meldrum BS (2000) Anticonvulsant activity of two metabotropic glutamate Group I antagonists selective for the mGlu5 receptor: 2-methyl-6-(phenylethynyl)-pyridine (MPEP), and (E)-6-methyl-2-styryl-pyridine (SIB 1893). *Neuropharmacology* 39:1567-1574.
- Charpak S, Gähwiler BH, Do KQ, Knöpfel T (1990) Potassium conductances in hippocampal neurons blocked by excitatory amino-acid transmitters. *Nature* 347:765.
- Charpak S, Gähwiler BH (1991) Glutamate mediates a slow synaptic response in hippocampal slice cultures. *Proceedings of the Royal Society B: Biological Sciences* 243:221-226.
- Chavis P, Shinozaki H, Bockaert J, Fagni L (1994) The metabotropic glutamate receptor types 2/3 inhibit L-type calcium channels via a pertussis toxin-sensitive G-protein in cultured cerebellar granule cells. *J Neurosci* 14:7067-7076.

- Chiamulera C, Epping-Jordan MP, Zocchi A, Marcon C, Cottiny Cc, Tacconi S, Corsi M, Orzi F, Conquet Fo (2001) Reinforcing and locomotor stimulant effects of cocaine are absent in mGluR5 null mutant mice. *Nat Neurosci* 4:873-874.
- Chittajallu R, Braithwaite SP, Clarke VRJ, Henley JM (1999) Kainate receptors: subunits, synaptic localization and function. *Trends in Pharmacological Sciences* 20:26-35.
- Cho RW, Park JM, Wolff SBE, Xu D, Hopf C, Kim J-a, Reddy RC, Petralia RS, Perin MS, Linden DJ, Worley PF (2008) mGluR1/5-Dependent Long-Term Depression Requires the Regulated Ectodomain Cleavage of Neuronal Pentraxin NPR by TACE. *Neuron* 57:858-871.
- Chu Z, Moenter SM (2005) Endogenous activation of metabotropic glutamate receptors modulates GABAergic transmission to gonadotropin-releasing hormone neurons and alters their firing rate: a possible local feedback circuit. *J Neurosci* 25:5740-5749.
- Clements JD (1996) Transmitter timecourse in the synaptic cleft: its role in central synaptic function. *Trends in Neurosciences* 19:163-171.
- Cobb SR, Buhl EH, Halasy K, Paulsen O, Somogyi P (1995) Synchronization of neuronal activity in hippocampus by individual GABAergic interneurons. *Nature* 378:75-78.
- Congar P, Leinekugel X, Ben-Ari Y, Crepel V (1997) A long-lasting calcium-activated nonselective cationic current is generated by synaptic stimulation or exogenous activation of group I metabotropic glutamate receptors in CA1 pyramidal neurons. *J Neurosci* 17:5366-5379.
- Conn PJ (2003) Physiological Roles and Therapeutic Potential of Metabotropic Glutamate Receptors. *Annals of the New York Academy of Sciences* 1003:12-21.
- Conn PJ, Battaglia G, Marino MJ, Nicoletti F (2005) Metabotropic glutamate receptors in the basal ganglia motor circuit. *Nat Rev Neurosci* 6:787-798.
- Conn PJ, Pin J-P (1997) Pharmacology and functions of metabotropic glutamate receptors. *Annual Review of Pharmacology and Toxicology* 37:205-237.

- Conquet F, Bashir ZI, Davies CH, Daniel H, Ferraguti F, Bordi F, Franz-Bacon K, Reggiani A, Matarrese V, Conde F, Collingridge GL, Crepel F (1994) Motor deficit and impairment of synaptic plasticity in mice lacking mGluR1. *Nature* 372:237-243.
- Contractor A, Gereau RW, Green T, Heinemann SF (1998) Direct effects of metabotropic glutamate receptor compounds on native and recombinant N-methyl-d-aspartate receptors. *Proceedings of the National Academy of Sciences of the United States of America* 95:8969-8974.
- Cooke SF, Bliss TVP (2006) Plasticity in the human central nervous system. *Brain* 129:1659-1673.
- Cotman CW, Monaghan DT (1988) Excitatory Amino Acid Neurotransmission: NMDA Receptors and Hebb-Type Synaptic Plasticity. *Annual Review of Neuroscience* 11:61-80.
- Coussen F (2009) Molecular determinants of kainate receptor trafficking. *Neuroscience* 158:25-35.
- Csicsvari J, Hirase H, Czurko A, Buzsáki G (1998) Reliability and state dependence of pyramidal cell-interneuron synapses in the hippocampus: an ensemble approach in the behaving rat. *Neuron* 21:179-189.
- Cull-Candy S, Brickley S, Farrant M (2001) NMDA receptor subunits: diversity, development and disease. *Current Opinion in Neurobiology* 11:327-335.
- Dale LB, Babwah AV, Ferguson SSG (2002) Mechanisms of metabotropic glutamate receptor desensitization: role in the patterning of effector enzyme activation. *Neurochemistry International* 41:319-326.
- Daw NW, Stein PSG, Fox K (1993) The Role of NMDA Receptors in Information Processing. *Annual Review of Neuroscience* 16:207-222.
- Daw MI, Bortolotto ZA, Saulle E, Zaman S, Collingridge GL, Isaac JTR (2002) Phosphatidylinositol 3 kinase regulates synapse specificity of hippocampal long-term depression. *Nat Neurosci* 5:835-836.
- De Blasi A, Conn PJ, Pin J-P, Nicoletti F (2001) Molecular determinants of metabotropic glutamate receptor signaling. *Trends in Pharmacological Sciences* 22:114-120.

- Debanne D, Gähwiler BH, Thompson SM (1996) Cooperative interactions in the induction of long-term potentiation and depression of synaptic excitation between hippocampal CA3-CA1 cell pairs in vitro. *Proceedings of the National Academy of Sciences of the United States of America* 93:11225-11230.
- Debanne D, Thompson SM (1996) Associative long-term depression in the hippocampus in vitro. *Hippocampus* 6:9-16.
- Debanne D, Daoudal G, Sourdet V, Russier M (2003) Brain plasticity and ion channels. *Journal of Physiology-Paris* 97:403-414.
- Derkach VA, Oh MC, Guire ES, Soderling TR (2007) Regulatory mechanisms of AMPA receptors in synaptic plasticity. *Nat Rev Neurosci* 8:101-113.
- Desai MA, Conn PJ (1991) Excitatory effects of ACPD receptor activation in the hippocampus are mediated by direct effects on pyramidal cells and blockade of synaptic inhibition. *J Neurophysiol* 66:40-52.
- Dhami GK, Ferguson SSG (2006) Regulation of metabotropic glutamate receptor signaling, desensitization and endocytosis. *Pharmacology & Therapeutics* 111:260-271.
- Dingledine R, Borges K, Bowie D, Traynelis SF (1999) The Glutamate Receptor Ion Channels. *Pharmacol Rev* 51:7-62.
- Dragoi G, Carpi D, Recce M, Csicsvari J, Buzsaki G (1999) Interactions between hippocampus and medial septum during sharp waves and theta oscillation in the behaving rat. *J Neurosci* 19:6191-6199.
- Dudek SM, Bear MF (1992) Homosynaptic long-term depression in area CA1 of hippocampus and effects of N-methyl-D-aspartate receptor blockade. *Proceedings of the National Academy of Sciences of the United States of America* 89:4363-4367.
- Eaton SA, Birse EF, Wharton B, Sunter DC, Udvarhelyi PM, Watkins JC, Salt TE (1993) Mediation of thalamic sensory responses in vivo by ACPD-activated excitatory amino acid receptors. *European Journal of Neuroscience* 5:186-189.

- Enoki R, Hu Y-I, Hamilton D, Fine A (2009) Expression of long-term plasticity at individual synapses in hippocampus is graded, bidirectional, and mainly presynaptic: optical quantal analysis. *Neuron* 62:242-253.
- Esteban JA (2003) AMPA Receptor Trafficking: A Road Map for Synaptic Plasticity. *Mol Interv* 3:375-385.
- Fagni L, Ango F, Perroy J, Bockaert J (2004) Identification and functional roles of metabotropic glutamate receptor-interacting proteins. *Seminars in Cell & Developmental Biology* 15:289-298.
- Feinmark SJ, Begum R, Tsvetkov E, Goussakov I, Funk CD, Siegelbaum SA, Bolshakov VY (2003) 12-Lipoxygenase Metabolites of Arachidonic Acid Mediate Metabotropic Glutamate Receptor-Dependent Long-Term Depression at Hippocampal CA3-CA1 Synapses. *J Neurosci* 23:11427-11435.
- Ferraguti F, Conquet F, Corti C, Grandes P, Kuhn R, Knopfel T (1998) Immunohistochemical localization of the mGluR1beta metabotropic glutamate receptor in the adult rodent forebrain: Evidence for a differential distribution of mGluR1 splice variants. *The Journal of Comparative Neurology* 400:391-407.
- Ferraguti F, Cobden P, Pollard M, Cope D, Shigemoto R, Watanabe M, Somogyi P (2004) Immunolocalization of metabotropic glutamate receptor 1alpha (mGluR1alpha) in distinct classes of interneuron in the CA1 region of the rat hippocampus. *Hippocampus* 14:193-215.
- Ferraguti F, Shigemoto R (2006) Metabotropic glutamate receptors. *Cell and Tissue Research* 326:483-504.
- Finch EA, Augustine GJ (1998) Local calcium signalling by inositol-1,4,5-trisphosphate in Purkinje cell dendrites. *Nature* 396:753-756.
- Fiorillo CD, Williams JT (1998) Glutamate mediates an inhibitory postsynaptic potential in dopamine neurons. *Nature* 394:78-82.
- Fischer Y, Gahwiler BH, Thompson SM (1999) Activation of intrinsic hippocampal theta oscillations by acetylcholine in rat septo-hippocampal cocultures. *J Physiol* 519:405-413.

- Fischer Y, Wittner L, Freund TF, Gähwiler BH (2002) Simultaneous activation of gamma and theta network oscillations in rat hippocampal slice cultures. *J Physiol* 539:857-868.
- Fitzjohn SM, Kingston AE, Lodge D, Collingridge GL (1999) DHPG-induced LTD in area CA1 of juvenile rat hippocampus; characterisation and sensitivity to novel mGlu receptor antagonists. *Neuropharmacology* 38:1577-1583.
- Fitzjohn SM, Palmer MJ, May JE, Neeson A, Morris SA, Collingridge GL (2001) A characterisation of long-term depression induced by metabotropic glutamate receptor activation in the rat hippocampus in vitro. *J Physiol (Lond)* 537:421-430.
- Flint AC, Dammerman RS, Kriegstein AR (1999) Endogenous activation of metabotropic glutamate receptors in neocortical development causes neuronal calcium oscillations. *Proceedings of the National Academy of Sciences of the United States of America* 96:12144-12149.
- Fotuhi M, Sharp AH, Glatt CE, Hwang PM, von Krosigk M, Snyder SH, Dawson TM (1993) Differential localization of phosphoinositide-linked metabotropic glutamate receptor (mGluR1) and the inositol 1,4,5-trisphosphate receptor in rat brain. *J Neurosci* 13:2001-2012.
- Freund TF, Antal M (1988) GABA-containing neurons in the septum control inhibitory interneurons in the hippocampus. *Nature* 336:170-173.
- Gähwiler BH, Thompson SM, McKinney RA, Debanne D, Robertsonm RT (1998) Culturing nerve cells. In: *Organotypic slice cultures of neural tissue* (Banker G, Goslin K, eds), pp461-498. Cambridge, MA: MIT.
- Gallagher SM, Daly CA, Bear MF, Huber KM (2004) Extracellular Signal-Regulated Protein Kinase Activation Is Required for Metabotropic Glutamate Receptor-Dependent Long-Term Depression in Hippocampal Area CA1. *J Neurosci* 24:4859-4864.
- Gee CE, Benquet P, Gerber U (2003) Group I metabotropic glutamate receptors activate a calcium-sensitive transient receptor potential-like conductance in rat hippocampus. *J Physiol (Lond)* 546:655-664.

- Gee CE, Lacaille JC (2004) Group I metabotropic glutamate receptor actions in oriens/alveus interneurons of rat hippocampal CA1 region. *Brain Research* 1000:92-101.
- Gerber U (2002) G-protein-coupled receptors, tyrosine kinases and neurotransmission. *Neuropharmacology* 42:587-592.
- Gerber U, Gee CE, Benquet P (2007) Metabotropic glutamate receptors: intracellular signaling pathways. *Current Opinion in Pharmacology* 7:56-61.
- Gereau RW, Conn PJ (1995a) Multiple presynaptic metabotropic glutamate receptors modulate excitatory and inhibitory synaptic transmission in hippocampal area CA1. *J Neurosci* 15:6879.
- Gereau RW, Conn PJ (1995b) Roles of specific metabotropic glutamate receptor subtypes in regulation of hippocampal CA1 pyramidal cell excitability. *J Neurophysiol* 74:122-129.
- Geurts JJG, Wolswijk G, Bo L, van der Valk P, Polman CH, Troost D, Aronica E (2003) Altered expression patterns of group I and II metabotropic glutamate receptors in multiple sclerosis. *Brain* 126:1755-1766.
- Glaum SR, Miller RJ (1992) Metabotropic glutamate receptors mediate excitatory transmission in the nucleus of the solitary tract. *J Neurosci* 12:2251-2258.
- Godwin DW, Van Horn SC, Sesma M, Romano C, Sherman SM (1996) Ultrastructural localization suggests that retinal and cortical inputs access different metabotropic glutamate receptors in the lateral geniculate nucleus. *J Neurosci* 16:8181-8192.
- Göres T, Penke B, Böti Z, Katarova Z, Hámori J (1993) Immunohistochemical visualization of a metabotropic glutamate receptor. *NeuroReport* 4:283-286.
- Grastyán E, Lissák K, Madarász I, Donhoffer H (1959) Hippocampal electrical activity during the development of conditioned reflexes. *Electroencephalography and Clinical Neurophysiology* 11:409-430.
- Green JD, Arduini AA (1954) Hippocampal electrical activity in arousal. *J Neurophysiol* 17:533-557.

- Greger IH, Esteban JA (2007) AMPA receptor biogenesis and trafficking. *Current Opinion in Neurobiology* 17:289-297.
- Greger IH, Ziff EB, Penn AC (2007) Molecular determinants of AMPA receptor subunit assembly. *Trends in Neurosciences* 30:407-416.
- Grenier F, Timofeev I, Steriade M (2001) Focal Synchronization of Ripples (80-200 Hz) in Neocortex and Their Neuronal Correlates. *J Neurophysiol* 86:1884-1898.
- Grenier F, Timofeev I, Steriade M (2003) Neocortical Very Fast Oscillations (Ripples, 80-200 Hz) During Seizures: Intracellular Correlates. *J Neurophysiol* 89:841-852.
- Grishin AA, Gee CE, Gerber U, Benquet P (2004) Differential calcium-dependent modulation of NMDA currents in CA1 and CA3 hippocampal pyramidal cells. *J Neurosci* 24:350-355.
- Groc L, Choquet D (2006) AMPA and NMDA glutamate receptor trafficking: multiple roads for reaching and leaving the synapse. *Cell and Tissue Research* 326:423-438.
- Grueter B, McElligott Z, Winder D (2007) Group I mGluRs and Long-Term Depression: Potential Roles in Addiction? *Molecular Neurobiology* 36:232-244.
- Gubellini P, Saulle E, Centonze D, Bonsi P, Pisani A, Bernardi G, Conquet F, Calabresi P (2001) Selective involvement of mGlu1 receptors in corticostriatal LTD. *Neuropharmacology* 40:839-846.
- Gubellini P, Saulle E, Centonze D, Costa C, Tropepi D, Bernardi G, Conquet F, Calabresi P (2003) Corticostriatal LTP requires combined mGluR1 and mGluR5 activation. *Neuropharmacology* 44:8-16.
- Guérineau NC, Gähwiler BH, Gerber U (1994) Reduction of resting K⁺ current by metabotropic glutamate and muscarinic receptors in rat CA3 cells: mediation by G-proteins. *J Physiol* 474:27-33.
- Guérineau NC, Bossu JL, Gähwiler BH, Gerber U (1995) Activation of a nonselective cationic conductance by metabotropic glutamatergic and

- muscarinic agonists in CA3 pyramidal neurons of the rat hippocampus. *J Neurosci* 15:4395-4407.
- Guo W, Wei F, Zou S, Robbins MT, Sugiyo S, Ikeda T, Tu J-C, Worley PF, Dubner R, Ren K (2004) Group I metabotropic glutamate receptor NMDA receptor coupling and signaling cascade mediate spinal dorsal horn NMDA receptor 2B tyrosine phosphorylation associated with inflammatory hyperalgesia. *J Neurosci* 24:9161-9173.
- Gulyás AI, Hájos N, Katona I, Freund TF (2003) Interneurons are the local targets of hippocampal inhibitory cells which project to the medial septum. *European Journal of Neuroscience* 17:1861-1872.
- Halasy K, Buhl EH, Lörinczi Z, Tamás G, Somogyi P (1996) Synaptic target selectivity and input of GABAergic basket and bistratified interneurons in the CA1 area of the rat hippocampus. *Hippocampus* 6:306-329.
- Hannan AJ, Blakemore C, Katsnelson A, Vitalis T, Huber KM, Bear M, Roder J, Kim D, Shin H-S, Kind PC (2001) PLC- β 1, activated via mGluRs, mediates activity-dependent differentiation in cerebral cortex. *Nat Neurosci* 4:282-288.
- Hanson JE, Smith Y (1999) Group I metabotropic glutamate receptors at GABAergic synapses in monkeys. *J Neurosci* 19:6488-6496.
- Harney SC, Rowan M, Anwyl R (2006) Long-term depression of NMDA receptor-mediated synaptic transmission is dependent on activation of metabotropic glutamate receptors and is altered to long-term potentiation by low intracellular calcium buffering. *J Neurosci* 26:1128-1132.
- Harris KD, Hirase H, Leinekugel X, Henze DA, Buzsáki G (2001) Temporal interaction between single spikes and complex spike bursts in hippocampal pyramidal cells. *Neuron* 32:141-149.
- Harvey J, Collingridge GL (1993) Signal transduction pathways involved in the acute potentiation of NMDA responses by 1S,3R-ACPD in rat hippocampal slices. *Br J Pharmacol* 109:1085-1090.

- Hayashi Y, Momiyama A, Takahashi T, Ohishi H, Ogawa-Meguro R (1993) Role of a metabotropic glutamate receptor in synaptic modulation in the accessory olfactory bulb. *Nature* 366:687.
- Hensch TK, Stryker MP (1996) Ocular dominance plasticity under metabotropic glutamate receptor blockade. *Science* 272:554-557.
- Herman MA, Jahr CE (2007) Extracellular glutamate concentration in hippocampal slice. *J Neurosci* 27:9736-9741.
- Hermans E, Challiss RA (2001) Structural, signalling and regulatory properties of the group I metabotropic glutamate receptors: prototypic family C G-protein-coupled receptors. *Biochem J* 359:465-484.
- Herrero I, Miras-Portugal MT, Sanchez-Prieto J (1992) Positive feedback of glutamate exocytosis by metabotropic presynaptic receptor stimulation. *Nature* 360:163-166.
- Heuss C, Scanziani M, Gahwiler BH, Gerber U (1999) G-protein-independent signaling mediated by metabotropic glutamate receptors. *Nat Neurosci* 2:1070-1077.
- Hirose M, Kitano J, Nakajima Y, Moriyoshi K, Yanagi S, Yamamura H, Muto T, Jingami H, Nakanishi S (2004) Phosphorylation and Recruitment of Syk by Immunoreceptor Tyrosine-based activation motif-based phosphorylation of tamalin. *J Biol Chem* 279:32308-32315.
- Holbro N, Grunditz A, Oertner TG (2009) Differential distribution of endoplasmic reticulum controls metabotropic signaling and plasticity at hippocampal synapses. *Proceedings of the National Academy of Sciences* 106:15055-15060.
- Hollmann M, Heinemann S (1994) Cloned glutamate receptors. *Annual Review of Neuroscience* 17:31-108.
- Horn R, Marty A (1988) Muscarinic activation of ionic currents measured by a new whole-cell recording method. *J Gen Physiol* 92:145-159.
- Hsia AY, Malenka RC, Nicoll RA (1998) Development of excitatory circuitry in the hippocampus. *J Neurophysiol* 79:2013-2024.

- Huber KM, Kayser MS, Bear MF (2000) Role for Rapid Dendritic Protein Synthesis in Hippocampal mGluR-Dependent Long-Term Depression. *Science* 288:1254-1256.
- Huber KM, Roder JC, Bear MF (2001) Chemical Induction of mGluR5- and Protein Synthesis-Dependent Long-Term Depression in Hippocampal Area CA1. *J Neurophysiol* 86:321-325.
- Hubert GW, Paquet M, Smith Y (2001) Differential subcellular localization of mGluR1a and mGluR5 in the rat and monkey substantia nigra. *J Neurosci* 21:1838-1847.
- Hubert GW, Smith Y (2004) Age-related changes in the expression of axonal and glial group I metabotropic glutamate receptor in the rat substantia nigra pars reticulata. *The Journal of Comparative Neurology* 475:95-106.
- Huettnner JE (2003) Kainate receptors and synaptic transmission. *Progress in Neurobiology* 70:387-407.
- Ireland DR, Abraham WC (2009) Mechanisms of Group I mGluR-Dependent Long-Term Depression of NMDA Receptor-Mediated Transmission at Schaffer Collateral-CA1 Synapses. *J Neurophysiol* 101:1375-1385.
- Isaac JTR, Oliet SHR, Hjelmstad GO, Nicoll RA, Malenka RC (1996) Expression mechanisms of long-term potentiation in the hippocampus. *Journal of Physiology-Paris* 90:299-303.
- Isaac JTR, Mellor J, Hurtado D, Roche KW (2004) Kainate receptor trafficking: physiological roles and molecular mechanisms. *Pharmacology & Therapeutics* 104:163-172.
- Ishikawa K, Nash SR, Nishimune A, Neki A, Kaneko S, Nakanishi S (1999) Competitive interaction of Seven in absentia homolog-1A and Ca²⁺/calmodulin with the cytoplasmic tail of group 1 metabotropic glutamate receptors. *Genes to Cells* 4:381-390.
- Ito M (1989) Long-Term Depression. *Annual Review of Neuroscience* 12:85-102.
- Jaarsma D, Dino MR, Ohishi H, Shigemoto R, Mugnaini E (1998) Metabotropic glutamate receptors are associated with non-synaptic appendages of

- unipolar brush cells in rat cerebellar cortex and cochlear nuclear complex. *Journal of Neurocytology* 27:303-327.
- Jabaudon D, Shimamoto K, Yasuda-Kamatani Y, Scanziani M, Gähwiler BH, Gerber U (1999) Inhibition of uptake unmasks rapid extracellular turnover of glutamate of nonvesicular origin. *Proceedings of the National Academy of Sciences* 96:8733-8738.
- Jaskolski F, Coussen F, Mulle C (2005) Subcellular localization and trafficking of kainate receptors. *Trends in Pharmacological Sciences* 26:20-26.
- Jia H, Rustioni A, Valtschanoff J, G. (1999) Metabotropic glutamate receptors in superficial laminae of the rat dorsal horn. *The Journal of Comparative Neurology* 410:627-642.
- Jingami H, Nakanishi S, Morikawa K (2003) Structure of the metabotropic glutamate receptor. *Current Opinion in Neurobiology* 13:271-278.
- Jo J, Heon S, Kim MJ, Son GH, Park Y, Henley JM, Weiss JL, Sheng M, Collingridge GL, Cho K (2008) Metabotropic glutamate receptor-mediated LTD involves two interacting Ca²⁺ sensors, NCS-1 and PICK1. *Neuron* 60:1095-1111.
- Johnson JL (1978) The excitant amino acids glutamic and aspartic acid as transmitter candidates in the vertebrate central nervous system. *Progress in Neurobiology* 10:155-202.
- Johnston D, Williams S, Jaffe D, Gray R (1992) NMDA-Receptor-Independent Long-Term Potentiation. *Annual Review of Physiology* 54:489-505.
- Joly C, Gomeza J, Brabet I, Curry K, Bockaert J, Pin JP (1995) Molecular, functional, and pharmacological characterization of the metabotropic glutamate receptor type 5 splice variants: comparison with mGluR1. *J Neurosci* 15:3970-3981.
- Jonas P, Sakmann B (1992) Glutamate receptor channels in isolated patches from CA1 and CA3 pyramidal cells of rat hippocampal slices. *The Journal of Physiology* 455:143-171.
- Jouvet M (1969) Biogenic amines and the states of sleep. *Science* 163:32-41.

- Kamiya H (2002) Kainate receptor-dependent presynaptic modulation and plasticity. *Neuroscience Research* 42:1-6.
- Kammermeier PJ, Xiao B, Tu JC, Worley PF, Ikeda SR (2000) Homer proteins regulate coupling of group I metabotropic glutamate receptors to N-type calcium and M-type potassium channels. *J Neurosci* 20:7238-7245.
- Kammermeier P, Ikeda S (2001) A role for Seven in Absentia Homolog (Siah1a) in metabotropic glutamate receptor signaling. *BMC Neuroscience* 2:15.
- Kamondi A, Acsády L, Wang X-J, Buzsáki G (1998) Theta oscillations in somata and dendrites of hippocampal pyramidal cells in vivo: Activity-dependent phase-precession of action potentials. *Hippocampus* 8:244-261.
- Kano M, Hashimoto K, Kurihara H, Watanabe M, Inoue Y, Aiba A, Tonegawa S (1997) Persistent multiple climbing fiber innervation of cerebellar purkinje cells in mice lacking mGluR1. *Neuron* 18:71-79.
- Karim F, Bhawe G, Gereau RW (2001) Metabotropic glutamate receptors on peripheral sensory neuron terminals as targets for the development of novel analgesics. *Molecular Psychiatry* 6:615-617.
- Kato N (1993) Dependence of long-term depression on postsynaptic metabotropic glutamate receptors in visual cortex. *Proceedings of the National Academy of Sciences of the United States of America* 90:3650-3654.
- Kato A, Ozawa F, Saitoh Y, Hirai K, Inokuchi K (1997) vesl, a gene encoding VASP/Ena family related protein, is upregulated during seizure, long-term potentiation and synaptogenesis. *FEBS Letters* 412:183-189.
- Kato A, Ozawa F, Saitoh Y, Fukazawa Y, Sugiyama H, Inokuchi K (1998) Novel members of the Vesl/Homer family of PDZ proteins that bind metabotropic glutamate receptors. *J Biol Chem* 273:23969-23975.
- Kemp N, Bashir ZI (1997) NMDA Receptor-dependent and -independent Long-term Depression in the CA1 Region of the Adult Rat Hippocampus In Vitro. *Neuropharmacology* 36:397-399.

- Kemp N, Bashir ZI (1999) Induction of LTD in the adult hippocampus by the synaptic activation of AMPA/kainate and metabotropic glutamate receptors. *Neuropharmacology* 38:495-504.
- Kemp N, Bashir ZI (2001) Long-term depression: a cascade of induction and expression mechanisms. *Progress in Neurobiology* 65:339-365.
- Kew JNC, Kemp JA (2005) Ionotropic and metabotropic glutamate receptor structure and pharmacology. *Psychopharmacology* 179:4-29.
- Kerchner GA, Nicoll RA (2008) Silent synapses and the emergence of a postsynaptic mechanism for LTP. *Nat Rev Neurosci* 9:813-825.
- Kerner JA, Standaert DG, Penney JB, Young AB, Landwehrmeyer GB (1997) Expression of group one metabotropic glutamate receptor subunit mRNAs in neurochemically identified neurons in the rat neostriatum, neocortex, and hippocampus. *Molecular Brain Research* 48:259-269.
- Kessels HW, Malinow R (2009) Synaptic AMPA Receptor Plasticity and Behavior. *Neuron* 61:340-350.
- King C, Recce M, Keefe JO (1998) The rhythmicity of cells of the medial septum/diagonal band of Broca in the awake freely moving rat: relationships with behaviour and hippocampal theta. *European Journal of Neuroscience* 10:464-477.
- Kingston AE, O'Neill MJ, Bond A, Bruno V, Battaglia G, Nicoletti F, Harris JR, Clark BP, Monn JA, Lodge D, Schoepp DD (1999) Neuroprotective actions of novel and potent ligands of group I and group II metabotropic glutamate receptors. *Annals of the New York Academy of Sciences* 890:438-449.
- Kitano J, Kimura K, Yamazaki Y, Soda T, Shigemoto R, Nakajima Y, Nakanishi S (2002) Tamalin, a PDZ domain-containing protein, links a protein complex formation of group 1 metabotropic glutamate receptors and the guanine nucleotide exchange factor cytohesins. *J Neurosci* 22:1280-1289.
- Köhr G (2006) NMDA receptor function: subunit composition versus spatial distribution. *Cell and Tissue Research* 326:439-446.

- Konopacki J, Bruce MacIver M, Bland BH, Roth SH (1987) Carbachol-induced EEG 'theta' activity in hippocampal brain slices. *Brain Research* 405:196-198.
- Krause M, Offermanns S, Stocker M, Pedarzani P (2002) Functional specificity of α_q and α_{11} in the cholinergic and glutamatergic modulation of potassium currents and excitability in hippocampal neurons. *J Neurosci* 22:666-673.
- Kullmann DM, Siegelbaum SA (1995) The site of expression of NMDA receptor-dependent LTP: New fuel for an old fire. *Neuron* 15:997-1002.
- Kullmann DM (2001) Presynaptic Kainate Receptors in the Hippocampus: Slowly Emerging from Obscurity. *Neuron* 32:561-564.
- Kullmann DM, Lamsa KP (2007) Long-term synaptic plasticity in hippocampal interneurons. *Nat Rev Neurosci* 8:687-699.
- Lan J-y, Skeberdis VA, Jover T, Zheng X, Bennett MVL, Zukin RS (2001) Activation of metabotropic glutamate receptor 1 accelerates NMDA receptor trafficking. *J Neurosci* 21:6058-6068.
- Laslo P, Lipski J, Funk GD (2001) Differential expression of Group I metabotropic glutamate receptors in motoneurons at low and high risk for degeneration in ALS. *NeuroReport* 12:1903-1908.
- Lau CG, Zukin RS (2007) NMDA receptor trafficking in synaptic plasticity and neuropsychiatric disorders. *Nat Rev Neurosci* 8:413-426.
- Lawson VH, Bland BH (1993) The Role of the Septohippocampal Pathway in the regulation of hippocampal field activity and behavior: analysis by the intraseptal microinfusion of carbachol, atropine, and procaine. *Experimental Neurology* 120:132-144.
- Lerma J (2003) Roles and rules of kainate receptors in synaptic transmission. *Nat Rev Neurosci* 4:481-495.
- Lerma J (2006) Kainate receptor physiology. *Current Opinion in Pharmacology* 6:89-97.
- Levy LM, Warr O, Attwell D (1998) Stoichiometry of the glial glutamate transporter GLT-1 expressed inducibly in a chinese hamster ovary cell line

- selected for low endogenous Na⁺-dependent glutamate uptake. *J Neurosci* 18:9620-9628.
- Lisman JE (2009) The pre/post LTP debate. *Neuron* 63:281-284.
- Lin H-C, Wang S-J, Luo M-Z, Gean P-W (2000) Activation of group II metabotropic glutamate receptors induces long-term depression of synaptic transmission in the rat amygdala. *J Neurosci* 20:9017-9024.
- Linden DJ (1994) Long-term synaptic depression in the mammalian brain. *Neuron* 12:457-472.
- Linden DJ, Connor JA (1995) Long-Term Synaptic Depression. *Annual Review of Neuroscience* 18:319-357.
- Linden DJ, Dickinson MH, Smeyne M, Connor JA (1991) A long-term depression of AMPA currents in cultured cerebellar purkinje neurons. *Neuron* 7:81-89.
- Linden AM, Johnson BG, Peters SC, Shannon HE, Tian M, Wang Y, Yu JL, Köster A, Baez M, Schoepp DD (2002) Increased anxiety-related behavior in mice deficient for metabotropic glutamate 8 (mGlu8) receptor. *Neuropharmacology* 43:251-259.
- Lisman JE, Idiart MA (1995) Storage of 7 +/- 2 short-term memories in oscillatory subcycles. *Science* 267:1512-1515.
- Liu X-B, Muñoz A, Jones E, G (1998) Changes in subcellular localization of metabotropic glutamate receptor subtypes during postnatal development of mouse thalamus. *The Journal of Comparative Neurology* 395:450-465.
- Lopes da Silva FH, Witter MP, Boeijinga PH, Lohman AH (1990) Anatomic organization and physiology of the limbic cortex. *Physiol Rev* 70:453-511.
- Lopez-Bendito G, Shigemoto R, Fairen A, Lujan R (2002) Differential distribution of group I metabotropic glutamate receptors during rat cortical development. *Cereb Cortex* 12:625-638.
- Lu Y-M, Jia Z, Janus C, Henderson JT, Gerlai R, Wojtowicz JM, Roder JC (1997) Mice Lacking Metabotropic Glutamate Receptor 5 Show Impaired Learning and Reduced CA1 Long-Term Potentiation (LTP) But Normal CA3 LTP. *J Neurosci* 17:5196-5205.

- Lujan R, Nusser Z, Roberts JDB, Shigemoto R, Somogyi P (1996) Perisynaptic location of metabotropic glutamate receptors mGluR1 and mGluR5 on dendrites and dendritic spines in the rat hippocampus. *European Journal of Neuroscience* 8:1488-1500.
- Lujan R, Roberts JDB, Shigemoto R, Ohishi H, Somogyi P (1997) Differential plasma membrane distribution of metabotropic glutamate receptors mGluR1 α , mGluR2 and mGluR5, relative to neurotransmitter release sites. *Journal of Chemical Neuroanatomy* 13:219-241.
- Madison DV, Malenka RC, Nicoll RA (1991) Mechanisms Underlying Long-Term Potentiation of Synaptic Transmission. *Annual Review of Neuroscience* 14:379-397.
- Malenka RC (1994) Synaptic plasticity in the hippocampus: LTP and LTD. *Cell* 78:535-538.
- Malenka RC, Nicoll RA (1999) Long-Term Potentiation--A Decade of Progress? *Science* 285:1870-1874.
- Malenka RC (2003) Synaptic Plasticity and AMPA Receptor Trafficking. *Annals of the New York Academy of Sciences* 1003:1-11.
- Malenka RC, Bear MF (2004) LTP and LTD: An Embarrassment of Riches. *Neuron* 44:5-21.
- Malinow R, Malenka RC (2002) AMPA Receptor Trafficking and Synaptic Plasticity. *Annual Review of Neuroscience* 25:103-126.
- Manahan-Vaughan D (1997) Group 1 and 2 Metabotropic glutamate receptors play differential roles in hippocampal long-term depression and long-term potentiation in freely moving rats. *J Neurosci* 17:3303-3311.
- Mann EO, Paulsen O (2007) Role of GABAergic inhibition in hippocampal network oscillations. *Trends in Neurosciences* 30:343-349.
- Manzoni O, Bockaert J (1995) Metabotropic glutamate receptors inhibiting excitatory synapses in the CA1 area of rat hippocampus. *European Journal of Neuroscience* 7:2518-2523.

- Markowska AL, Olton DS, Givens B (1995) Cholinergic manipulations in the medial septal area: age-related effects on working memory and hippocampal electrophysiology. *J Neurosci* 15:2063-2073.
- Markram H, Lubke J, Frotscher M, Sakmann B (1997) Regulation of Synaptic Efficacy by Coincidence of Postsynaptic APs and EPSPs. *Science* 275:213-215.
- Martin LJ, Blackstone CD, Huganir RL, Price DL (1992) Cellular localization of a metabotropic glutamate receptor in rat brain. *Neuron* 9:259-270.
- Martinez JL, Derrick BE (1996) Long-term potentiation and learning. *Annual Review of Psychology* 47:173-203.
- Mateos JM, Azkue J, Benítez R, Sarría R, Losada J, Conquet F, Ferraguti F, Kuhn R, Knöpfel T, Grandes P (1998) Immunocytochemical localization of the mGluR1b metabotropic glutamate receptor in the rat hypothalamus. *The Journal of Comparative Neurology* 390:225-233.
- Mateos JM, Benítez R, Elezgarai I, Azkue JJ, Lázaro E, Osorio A, Bilbao A, Doñate F, Sarría R, Conquet F, Ferraguti F, Kuhn R, Knöpfel T, Grandes P (2000) Immunolocalization of the mGluR1b Splice Variant of the Metabotropic Glutamate Receptor 1 at Parallel Fiber-Purkinje Cell Synapses in the Rat Cerebellar Cortex. *Journal of Neurochemistry* 74:1301-1309.
- Matsuzaki M, Honkura N, Ellis-Davies GCR, Kasai H (2004) Structural basis of long-term potentiation in single dendritic spines. *Nature* 429:761-766.
- Matynia A, Kushner SA, Silva AJ (2002) Genetic Approaches to Molecular and Cellular Cognition: A Focus on LTP and Learning and Memory. *Annual Review of Genetics* 36:687-720.
- Mayer ML, Armstrong N (2004) Structure and Function of Glutamate Receptor Ion Channels. *Annual Review of Physiology* 66:161-181.
- Mayer ML (2005) Glutamate receptor ion channels. *Current Opinion in Neurobiology* 15:282-288.
- McCormick DA, von Krosigk M (1992) Corticothalamic activation modulates thalamic firing through glutamate "metabotropic" receptors. *Proceedings of*

- the National Academy of Sciences of the United States of America 89:2774-2778.
- McNaughton BL (1993) The Mechanism of Expression of Long-Term Enhancement of Hippocampal Synapses: Current Issues and Theoretical Implications. *Annual Review of Physiology* 55:375-396.
- Meldrum BS (2000) Glutamate as a Neurotransmitter in the Brain: Review of Physiology and Pathology. *J Nutr* 130:1007-1015.
- Meur KL, Galante M, Angulo MC, Audinat E (2007) Tonic activation of NMDA receptors by ambient glutamate of non-synaptic origin in the rat hippocampus. *J Physiol* 580:373-383.
- Miltner WHR, Braun C, ArnoldMatthias, Witte H, Taub E (1999) Coherence of gamma-band EEG activity as a basis for associative learning. *Nature* 397:434-436.
- Minakami R, Iida K-i, Hirakawa N (1995) The expression of two splice variants of metabotropic glutamate receptor subtype 5 in the rat brain and neuronal cells during development. *Journal of Neurochemistry* 65:1536-1542.
- Minakami R, Jinnai N, Sugiyama H (1997) Phosphorylation and calmodulin binding of the metabotropic glutamate receptor subtype 5 (mGluR5) are antagonistic in vitro. *J Biol Chem* 272:20291-20298.
- Miura M, Watanabe M, Offermanns S, Simon MI, Kano M (2002) Group I metabotropic glutamate receptor signaling via G α q/G α 11 secures the induction of long-term potentiation in the hippocampal area ca1. *J Neurosci* 22:8379-8390.
- Mori H, Mishina M (1995) Structure and function of the NMDA receptor channel. *Neuropharmacology* 34:1219-1237.
- Morsette DJ, Sidorowicz H, Sladek CD (2001) Role of metabotropic glutamate receptors in vasopressin and oxytocin release. *Am J Physiol Regul Integr Comp Physiol* 281:R452-458.
- Moult PR, Gladding CM, Sanderson TM, Fitzjohn SM, Bashir ZI, Molnar E, Collingridge GL (2006) Tyrosine Phosphatases Regulate AMPA Receptor

- Trafficking during Metabotropic Glutamate Receptor-Mediated Long-Term Depression. *J Neurosci* 26:2544-2554.
- Mulkey RM, Malenka RC (1992) Mechanisms underlying induction of homosynaptic long-term depression in area CA1 of the hippocampus. *Neuron* 9:967-975.
- Murthy VN, Fetz EE (1996) Oscillatory activity in sensorimotor cortex of awake monkeys: synchronization of local field potentials and relation to behavior. *J Neurophysiol* 76:3949-3967.
- Musante V, Neri E, Feligioni M, Puliti A, Pedrazzi M, Conti V, Usai C, Diaspro A, Ravazzolo R, Henley JM, Battaglia G, Pittaluga A (2008) Presynaptic mGlu1 and mGlu5 autoreceptors facilitate glutamate exocytosis from mouse cortical nerve endings. *Neuropharmacology* 55:474-482.
- Nahum-Levy R, Lipinski D, Shavit S, Benveniste M (2001) Desensitization of NMDA receptor channels is modulated by glutamate agonists. *Biophysical Journal* 80:2152-2166.
- Nahum-Levy R, Tam E, Shavit S, Benveniste M (2002) Glutamate but not glycine agonist affinity for NMDA receptors is influenced by small cations. *J Neurosci* 22:2550-2560.
- Nakamura T, Barbara J-G, Nakamura K, Ross WN (1999) Synergistic release of Ca²⁺ from IP₃-sensitive stores evoked by synaptic activation of mGluRs paired with backpropagating action potentials. *Neuron* 24:727-737.
- Nakanishi S (1994) Metabotropic glutamate receptors: Synaptic transmission, modulation, and plasticity. *Neuron* 13:1031-1037.
- Négyessy L, Vidnyánszky Z, Kuhn R, Knöpfel T, Görös TJ, Hátori J (1997) Light and electron microscopic demonstration of mGluR5 metabotropic glutamate receptor immunoreactive neuronal elements in the rat cerebellar cortex. *The Journal of Comparative Neurology* 385:641-650.
- Neki A, Ohishi H, Kaneko T, Shigemoto R, Nakanishi S, Mizuno N (1996) Metabotropic glutamate receptors mGluR2 and mGluR5 are expressed in two non-overlapping populations of Golgi cells in the rat cerebellum. *Neuroscience* 75:815-826.

- Netzeband JG, Parsons KL, Sweeney DD, Gruol DL (1997) Metabotropic glutamate receptor agonists alter neuronal excitability and Ca^{2+} levels via the phospholipase c transduction pathway in cultured purkinje neurons. *J Neurophysiol* 78:63-75.
- Neyman S, Manahan-Vaughan D (2008) Metabotropic glutamate receptor 1 (mGluR1) and 5 (mGluR5) regulate late phases of LTP and LTD in the hippocampal CA1 region *in vitro*. *European Journal of Neuroscience* 27:1345-1352.
- Nicoll RA, Oliet SHR, Malenka RC (1998) NMDA Receptor-Dependent and Metabotropic Glutamate Receptor-Dependent Forms of Long-Term Depression Coexist in CA1 Hippocampal Pyramidal Cells. *Neurobiology of Learning and Memory* 70:62-72.
- Nicoll RA, Tomita S, Brecht DS (2006) Auxiliary Subunits Assist AMPA-Type Glutamate Receptors. *Science* 311:1253-1256.
- Nusser Z, Mulvihill E, Streit P, Somogyi P (1994) Subsynaptic segregation of metabotropic and ionotropic glutamate receptors as revealed by immunogold localization. *Neuroscience* 61:421-427.
- O'Connor JJ, Rowan MJ, Anwyl R (1995) Tetanically induced LTP involves a similar increase in the AMPA and NMDA receptor components of the excitatory postsynaptic current: investigations of the involvement of mGlu receptors. *J Neurosci* 15:2013-2020.
- Oliet SHR, Malenka RC, Nicoll RA (1997) Two distinct forms of long-term depression coexist in CA1 hippocampal pyramidal cells. *Neuron* 18:969-982.
- Ong WY, He Y, Tan KK, Garey LJ (1998) Differential localisation of the metabotropic glutamate receptor mGluR1a and the ionotropic glutamate receptor GluR2/3 in neurons of the human cerebral cortex. *Experimental Brain Research* 119:367-374.
- Otani S, Daniel H, Takita M, Crepel F (2002) Long-term depression induced by postsynaptic group II metabotropic glutamate receptors linked to

- phospholipase c and intracellular calcium rises in rat prefrontal cortex. *J Neurosci* 22:3434-3444.
- Packard MG, Vecchioli SF, Schroeder JP, Gasbarri A (2001) Task-dependent role for dorsal striatum metabotropic glutamate receptors in memory. *Learning & Memory* 8:96-103.
- Palmer MJ, Irving AJ, Seabrook GR, Jane DE, Collingridge GL (1997) The group I mGlu receptor agonist DHPG induces a novel form of LTD in the CA1 region of the hippocampus. *Neuropharmacology* 36:1517-1532.
- Paoletti P, Neyton J (2007) NMDA receptor subunits: function and pharmacology. *Current Opinion in Pharmacology* 7:39-47.
- Park YK, Galik J, Ryu PD, Randic M (2004) Activation of presynaptic group I metabotropic glutamate receptors enhances glutamate release in the rat spinal cord substantia gelatinosa. *Neuroscience Letters* 361:220-224.
- Patneau DK, Mayer ML (1990) Structure-activity relationships for amino acid transmitter candidates acting at N-methyl-D-aspartate and quisqualate receptors. *J Neurosci* 10:2385-2399.
- Petralia RS, Wang YX, Zhao HM, Wenthold RJ (1996) Ionotropic and metabotropic glutamate receptors show unique postsynaptic, presynaptic, and glial localizations in the dorsal cochlear nucleus. *The Journal of Comparative Neurology* 372:356-383.
- Petsche H, Stumpf C, Gogolak G (1962) The significance of the rabbit's septum as a relay station between the midbrain and the hippocampus I. The control of hippocampus arousal activity by the septum cells. *Electroencephalography and Clinical Neurophysiology* 14:202-211.
- Phillis JW, Smith-Barbour M, Perkins LM, O'Regan MH (1994) Characterization of glutamate, aspartate, and GABA release from ischemic rat cerebral cortex. *Brain Research Bulletin* 34:457-466.
- Pin JP, Bockaert J (1995) Get receptive to metabotropic glutamate receptors. *Current Opinion in Neurobiology* 5:342-349.
- Pin JP, Duvoisin R (1995) The metabotropic glutamate receptors: structure and functions. *Neuropharmacology* 34:1.

- Pinheiro P, Mulle C (2006) Kainate receptors. *Cell and Tissue Research* 326:457-482.
- Pinheiro PS, Mulle C (2008) Presynaptic glutamate receptors: physiological functions and mechanisms of action. *Nat Rev Neurosci* 9:423-436.
- Pisani A, Calabresi P, Centonze D, Bernardi G (1997) Enhancement of NMDA responses by group I metabotropic glutamate receptor activation in striatal neurones. *British Journal of Pharmacology* 120:1007-1014.
- Pisani A, Gubellini P, Bonsi P, Conquet F, Picconi B, Centonze D, Bernardi G, Calabresi P (2001) Metabotropic glutamate receptor 5 mediates the potentiation of N-methyl--aspartate responses in medium spiny striatal neurons. *Neuroscience* 106:579-587.
- Plenz D, Kitai ST (1998) Regulation of the nigrostriatal pathway by metabotropic glutamate receptors during development. *J Neurosci* 18:4133-4144.
- Poli A, Beraudi A, Villani L, Storto M, Battaglia G, Di Giorgi Gerevini V, Cappuccio I, Caricasole A, D'Onofrio M, Nicoletti F (2003) Group II metabotropic glutamate receptors regulate the vulnerability to hypoxic brain damage. *J Neurosci* 23:6023-6029.
- Racca C, Stephenson FA, Streit P, Roberts JDB, Somogyi P (2000) NMDA receptor content of synapses in stratum radiatum of the hippocampal CA1 area. *J Neurosci* 20:2512-2522.
- Raghavachari S, Kahana MJ, Rizzuto DS, Caplan JB, Kirschen MP, Bourgeois B, Madsen JR, Lisman JE (2001) Gating of human theta oscillations by a working memory task. *J Neurosci* 21:3175-3183.
- Raymond CR, Thompson VL, Tate WP, Abraham WC (2000) Metabotropic glutamate receptors trigger homosynaptic protein synthesis to prolong long-term potentiation. *J Neurosci* 20:969-976.
- Redman S (1996) The hippocampus, long-term potentiation and memory. *Clinical and Experimental Pharmacology and Physiology* 23:961-964.
- Riedel G, Platt B, Micheau J (2003) Glutamate receptor function in learning and memory. *Behavioural Brain Research* 140:1-47.

- Rodriguez E, George N, Lachaux J-P, Martinerie J, Renault B, Varela FJ (1999) Perception's shadow: long-distance synchronization of human brain activity. *Nature* 397:430-433.
- Rodríguez-Moreno A, Sistiaga A, Lerma J, Sánchez-Prieto J (1998) Switch from facilitation to inhibition of excitatory synaptic transmission by group I mGluR desensitization. *Neuron* 21:1477-1486.
- Romano C, Sesma MA, McDonald CT, O'Malley K, Van den Pol AN (1995) Distribution of metabotropic glutamate receptor mGluR5 immunoreactivity in rat brain. *J Comp Neurol* 355:455.
- Romano C, van den Pol AN, O'Malley KL (1996) Enhanced early developmental expression of the metabotropic glutamate receptor mGluR5 in rat brain: Protein, mRNA splice variants, and regional distribution. *The Journal of Comparative Neurology* 367:403-412.
- Sah P, Hestrin S, Nicoll RA (1989) Tonic activation of NMDA receptors by ambient glutamate enhances excitability of neurons. *Science* 246:815-818.
- Sahara Y, Westbrook GL (1993) Modulation of calcium currents by a metabotropic glutamate receptor involves fast and slow kinetic components in cultured hippocampal neurons. *J Neurosci* 13:3041.
- Sansig G, Bushell TJ, Clarke VRJ, Rozov A, Burnashev N, Portet C, Gasparini F, Schmutz M, Klebs K, Shigemoto R, Flor PJ, Kuhn R, Knoepfel T, Schroeder M, Hampson DR, Collett VJ, Zhang C, Duvoisin RM, Collingridge GL, van der Putten H (2001) Increased seizure susceptibility in mice lacking metabotropic glutamate receptor 7. *J Neurosci* 21:8734-8745.
- Santos SD, Carvalho AL, Caldeira MV, Duarte CB (2009) Regulation of AMPA receptors and synaptic plasticity. *Neuroscience* 158:105-125.
- Scanziani M (2000) GABA spillover activates postsynaptic GABAB receptors to control rhythmic hippocampal activity. *Neuron* 25:673-681.
- Schenk U, Matteoli M (2004) Presynaptic AMPA receptors: more than just ion channels? *Biology of the Cell* 96:257-260.

- Schoepp DD (2001) Unveiling the Functions of Presynaptic Metabotropic Glutamate Receptors in the Central Nervous System. *J Pharmacol Exp Ther* 299:12-20.
- Schrader LA, Tasker JG (1997) Presynaptic modulation by metabotropic glutamate receptors of excitatory and inhibitory synaptic inputs to hypothalamic magnocellular neurons. *J Neurophysiol* 77:527-536.
- Sheng M (2001) Molecular organization of the postsynaptic specialization. *Proceedings of the National Academy of Sciences of the United States of America* 98:7058-7061.
- Shigemoto R, Nakanishi S, Mizuno N (1992) Distribution of the mRNA for a metabotropic glutamate receptor (mGluR1) in the central nervous system: An in situ hybridization study in adult and developing rat. *The Journal of Comparative Neurology* 322:121-135.
- Shigemoto R, Nomura S, Ohishi H, Sugihara H, Nakanishi S, Mizuno N (1993) Immunohistochemical localization of a metabotropic glutamate receptor, mGluR5, in the rat brain. *Neuroscience Letters* 163:53-57.
- Shigemoto R, Abe T, Nomura S, Nakanishi S, Hirano T (1994) Antibodies inactivating mGluR1 metabotropic glutamate receptor block long-term depression in cultured Purkinje cells. *Neuron* 12:1245-1255.
- Shigemoto R, Kinoshita A, Wada E, Nomura S, Ohishi H, Takada M, Flor PJ, Neki A, Abe T, Nakanishi S, Mizuno N (1997) Differential presynaptic localization of metabotropic glutamate receptor subtypes in the rat hippocampus. *J Neurosci* 17:7503-7522.
- Shirasaki T, Harata N, Akaike N (1994) Metabotropic glutamate response in acutely dissociated hippocampal CA1 pyramidal neurones of the rat. *The Journal of Physiology* 475:439-453.
- Shu Y, Hasenstaub A, McCormick DA (2003) Turning on and off recurrent balanced cortical activity. *Nature* 423:288-293.
- Sik A, Ylinen A, Penttonen M, Buzsaki G (1994) Inhibitory CA1-CA3-hilar region feedback in the hippocampus. *Science* 265:1722-1724.

- Simkus CR, Stricker C (2002) The contribution of intracellular calcium stores to mEPSCs recorded in layer II neurones of rat barrel cortex. *The Journal of Physiology* 545:521-535.
- Singer W, Gray CM (1995) Visual Feature Integration and the Temporal Correlation Hypothesis. *Annual Review of Neuroscience* 18:555-586.
- Snyder EM, Philpot BD, Huber KM, Dong X, Fallon JR, Bear MF (2001) Internalization of ionotropic glutamate receptors in response to mGluR activation. *Nat Neurosci* 4:1079-1085.
- Song I, Huganir RL (2002) Regulation of AMPA receptors during synaptic plasticity. *Trends in Neurosciences* 25:578-588.
- Sorra KE, Harris KM (1993) Occurrence and three-dimensional structure of multiple synapses between individual radiatum axons and their target pyramidal cells in hippocampal area CA1. *J Neurosci* 13:3736-3748.
- Sourdet V, Russier M, Daoudal G, Ankri N, Debanne D (2003) Long-Term Enhancement of Neuronal Excitability and Temporal Fidelity Mediated by Metabotropic Glutamate Receptor Subtype 5. *J Neurosci* 23:10238-10248.
- Sprengel R (2006) Role of AMPA receptors in synaptic plasticity. *Cell and Tissue Research* 326:447-455.
- Stanton PK, Chattarji S, Sejnowski TJ (1991) 2-Amino-3-phosphonopropionic acid, an inhibitor of glutamate-stimulated phosphoinositide turnover, blocks induction of homosynaptic long-term depression, but not potentiation, in rat hippocampus. *Neuroscience Letters* 127:61-66.
- Stewart M, Fox SE (1990) Do septal neurons pace the hippocampal theta rhythm? *Trends in Neurosciences* 13:163-169.
- Swanson CJ, Bures M, Johnson MP, Linden A-M, Monn JA, Schoepp DD (2005) Metabotropic glutamate receptors as novel targets for anxiety and stress disorders. *Nat Rev Drug Discov* 4:131-144.
- Swartz KJ, Bean BP (1992) Inhibition of calcium channels in rat CA3 pyramidal neurons by a metabotropic glutamate receptor. *J Neurosci* 12:4358.
- Takechi H, Eilers J, Konnerth A (1998) A new class of synaptic response involving calcium release in dendritic spines. *Nature* 396:757-760.

- Takumi Y, Ramirez-Leon V, Laake P, Rinvik E, Ottersen OP (1999) Different modes of expression of AMPA and NMDA receptors in hippocampal synapses. *Nat Neurosci* 2:618-624.
- Teyler TJ, DiScenna P (1987) Long-Term Potentiation. *Annual Review of Neuroscience* 10:131-161.
- Thiels E, Xie X, Yeckel MF, Barrionuevo G, Berger TW (1996) NMDA Receptor-dependent LTD in different subfields of hippocampus in vivo and in vitro. *Hippocampus* 6:43-51.
- Thomson AM, Radpour S (1991) Excitatory connections between CA1 pyramidal cells revealed by spike triggered averaging in slices of rat hippocampus are partially NMDA receptor mediated. *Eur J Neurosci* 3:587-601.
- Tong G, Jahr CE (1994) Multivesicular release from excitatory synapses of cultured hippocampal neurons. *Neuron* 12:51-59.
- Topolnik L, Azzi M, Morin F, Kougioumoutzakis A, Lacaille J-C (2006) mGluR1/5 subtype-specific calcium signalling and induction of long-term potentiation in rat hippocampal oriens/alveus interneurons. *J Physiol* 575:115-131.
- Toth K, Borhegyi Z, Freund TF (1993) Postsynaptic targets of GABAergic hippocampal neurons in the medial septum-diagonal band of Broca complex. *J Neurosci* 13:3712-3724.
- Traub RD, Bibbig A, LeBeau FEN, Buhl EH, Whittington MA (2004) Cellular mechanisms of neuronal population oscillations in the hippocampus in vitro. *Annual Review of Neuroscience* 27:247-278.
- Tu JC, Xiao B, Yuan JP, Lanahan AA, Leoffert K, Li M, Linden DJ, Worley PF (1998) Homer binds a novel proline-rich motif and links group 1 metabotropic glutamate receptors with IP3 receptors. *Neuron* 21:717-726.
- Tu JC, Xiao B, Naisbitt S, Yuan JP, Petralia RS, Brakeman P, Doan A, Aakalu VK, Lanahan AA, Sheng M, Worley PF (1999) Coupling of mGluR/Homer and PSD-95 complexes by the Shank family of postsynaptic density proteins. *Neuron* 23:583-592.

- Ugolini A, Corsi M, Bordi F (1997) Potentiation of NMDA and AMPA responses by group I mGluR in spinal cord motoneurons. *Neuropharmacology* 36:1047-1055.
- Ugolini A, Corsi M, Bordi F (1999) Potentiation of NMDA and AMPA responses by the specific mGluR5 agonist CHPG in spinal cord motoneurons. *Neuropharmacology* 38:1569-1576.
- Valenti O, Conn PJ, Marino MJ (2002) Distinct physiological roles of the Gq-coupled metabotropic glutamate receptors co-expressed in the same neuronal populations. *Journal of Cellular Physiology* 191:125-137.
- van den Pol AN, Kogelman L, Ghosh P, Liljelund P, Blackstone C (1994) Developmental regulation of the hypothalamic metabotropic glutamate receptor mGluR1. *J Neurosci* 14:3816-3834.
- van den Pol AN, Romano C, Ghosh P (1995) Metabotropic glutamate receptor mGluR5 subcellular distribution and developmental expression in hypothalamus. *The Journal of Comparative Neurology* 362:134-150.
- Vanderwolf CH (1969) Hippocampal electrical activity and voluntary movement in the rat. *Electroencephalography and Clinical Neurophysiology* 26:407-418.
- Vertes RP, Kocsis B (1997) Brainstem-diencephalo-septohippocampal systems controlling the theta rhythm of the hippocampus. *Neuroscience* 81:893-926.
- Vidnyánszky Z, Hamori J, Négyessy L, Rüegg D, Knopfel T, Kuhn R, Görcs TJ (1994) Cellular, and subcellular localization of the mGluR5a metabotropic glutamate receptor in rat spinal cord. *NeuroReport* 6:209-213.
- Vidnyánszky Z, Görcs TJ, Négyessy L, Borostyánkői Z, Kuhn R, Knöpfel T, Hámori J (1996) Immunocytochemical visualization of the mGluR1 a metabotropic glutamate receptor at synapses of corticothalamic terminals originating from area 17 of the rat. *European Journal of Neuroscience* 8:1061-1071.
- Vinogradova OS (1995) Expression, control, and probable functional significance of the neuronal theta-rhythm. *Progress in Neurobiology* 45:523-583.

- Wahl F, Hardy AM, Plotkine M, Boulu R, Symon L (1994) Extracellular glutamate during focal cerebral ischaemia in rats: time course and calcium dependency. *Journal of Neurochemistry* 63:1003-1011.
- Wang X-F, Daw NW, Jin X-t (1998) The effect of ACPD on the responses to NMDA and AMPA varies with layer in slices of rat visual cortex. *Brain Research* 812:186-192.
- Wang J, Arora A, Yang L, Parekar N, Zhang G, Liu X, Choe E, Mao L (2005) Phosphorylation of AMPA receptors. *Molecular Neurobiology* 32:237-249.
- Wang W, Zhang Z, Shang J, Jiang Z-z, Wang S, Liu Y, Zhang L-y (2008) Activation of group I metabotropic glutamate receptors induces long-term depression in the hippocampal CA1 region of adult rats in vitro. *Neuroscience Research* 62:43-50.
- Watabe AM, Carlisle HJ, O'Dell TJ. (2002) Postsynaptic induction and presynaptic expression of group 1 mGluR-dependent LTD in the hippocampal CA1 region. *J Neurophysiol* 87:1395-1403.
- Waung MW, Pfeiffer BE, Nosyreva ED, Ronesi JA, Huber KM (2008) Rapid Translation of Arc/Arg3.1 Selectively Mediates mGluR-Dependent LTD through Persistent Increases in AMPAR Endocytosis Rate. *Neuron* 59:84-97.
- Wenthold RJ, Prybylowski K, Standley S, Sans N, Petralia RS (2003) Trafficking of NMDA Receptors. *Annual Review of Pharmacology and Toxicology* 43:335-358.
- White A-M, Kylänpää RA, Christie LA, McIntosh SJ, Irving AJ, Platt B (2003) Presynaptic group I metabotropic glutamate receptors modulate synaptic transmission in the rat superior colliculus via 4-AP sensitive K⁺ channels. *British Journal of Pharmacology* 140:1421-1433.
- Winson J (1978) Loss of hippocampal theta rhythm results in spatial memory deficit in the rat. *Science* 201:160-163.
- Wittmann M, Marino MJ, Bradley SR, Conn PJ (2001) Activation of group III mGluRs inhibits GABAergic and glutamatergic transmission in the substantia nigra pars reticulata. *J Neurophysiol* 85:1960-1968.

- Wu J, Rush A, Rowan MJ, Anwyl R (2001) NMDA receptor- and metabotropic glutamate receptor-dependent synaptic plasticity induced by high frequency stimulation in the rat dentate gyrus in vitro. *The Journal of Physiology* 533:745-755.
- Wu M, Hajszan T, Xu C, Leranth C, Alreja M (2004) Group I Metabotropic glutamate receptor activation produces a direct excitation of identified septohippocampal cholinergic neurons. *J Neurophysiol* 92:1216-1225.
- Xi Z-X, Shen H, Baker DA, Kalivas PW (2003) Inhibition of non-vesicular glutamate release by group III metabotropic glutamate receptors in the nucleus accumbens. *Journal of Neurochemistry* 87:1204-1212.
- Xiao B, Tu JC, Petralia RS, Yuan JP, Doan A, Breder CD, Ruggiero A, Lanahan AA, Wenthold RJ, Worley PF (1998) Homer regulates the association of group 1 metabotropic glutamate receptors with multivalent complexes of homer-related, synaptic proteins. *Neuron* 21:707-716.
- Xiao B, Cheng Tu J, Worley PF (2000) Homer: a link between neural activity and glutamate receptor function. *Current Opinion in Neurobiology* 10:370-374.
- Xiao M-Y, Zhou Q, Nicoll RA (2001) Metabotropic glutamate receptor activation causes a rapid redistribution of AMPA receptors. *Neuropharmacology* 41:664-671.
- Yu SP, Sensi SL, Canzoniero LM, Buisson A, Choi DW (1997) Membrane-delimited modulation of NMDA currents by metabotropic glutamate receptor subtypes 1/5 in cultured mouse cortical neurons. *The Journal of Physiology* 499:721-732.
- Zerangue N, Kavanaugh MP (1996) Flux coupling in a neuronal glutamate transporter. *Nature* 383:634-637.
- Zhang X-l, Zhou Z-y, Winterer J, Muller W, Stanton PK (2006) NMDA-dependent, but not group I metabotropic glutamate receptor-dependent, long-term depression at Schaffer collateral-CA1 synapses is associated with long-term reduction of release from the rapidly recycling presynaptic vesicle pool. *J Neurosci* 26:10270-10280.

- Zho W-M, You J-L, Huang C-C, Hsu K-S (2002) The Group I Metabotropic Glutamate Receptor Agonist (S)-3,5-Dihydroxyphenylglycine Induces a Novel Form of Depotentiation in the CA1 Region of the Hippocampus. *J Neurosci* 22:8838-8849.
- Zhong J, Gerber G, Kojic L, Randic M (2000) Dual modulation of excitatory synaptic transmission by agonists at group I metabotropic glutamate receptors in the rat spinal dorsal horn. *Brain Research* 887:359-377.
- Zhou S, Komak S, Du J, Carlton SM (2001) Metabotropic glutamate 1 α receptors on peripheral primary afferent fibers: their role in nociception. *Brain Research* 913:18-26.
- Zhou Q, Homma KJ, Poo MM (2004) Shrinkage of dendritic spines associated with long-term depression of hippocampal synapses. *Neuron* 44:749-757.
- Zimmer J, Gähwiler BH (1984) Cellular and connective organization of slice cultures of the rat hippocampus and fascia dentata. *J Comp Neurol* 228:432-446.

List of abbreviations

ACSF	artificial cerebrospinal fluid
AMPA	α -amino-3-hydroxy-5-methylisoxazole-4-propionic acid
AMPA	AMPA receptor
ATP	adenosine triphosphate
CaM	calmodulin
CaMKII	calcium/calmodulin-dependent protein kinase II
CF	climbing fiber
CNQX	6-cyano-7-nitroquinoxaline-2,3-dione
CNS	central nervous system
D-AP5	D-(-)-2-amino-5-phosphonopentanoic acid
DHPG	dihydroxyphenylglycine
EPSC	excitatory postsynaptic current
GABA	γ -aminobutyric acid
GPCR	G-protein-coupled receptor
GRIP	glutamate receptor interacting protein
GTP	guanosine triphosphate
iGluRs	ionotropic glutamate receptors
IN	interneuron
ING	interneuron network gamma
IP ₃	inositol-1,4,5-trisphosphate
KAR	kainate receptor
L-AP4	L-(+)-2-amino-4-phosphonobutyrate
LFS	low frequency stimulation
LTD	long-term depression
LTP	long-term potentiation
mAChR	muscarinic acetylcholine receptor
MCPG	α -methyl-4-carboxyphenylglycine
mGluR	metabotropic glutamate receptor
MPEP	2-methyl-6-(phenylethynyl)pyridine hydrochloride

MS-DBB	medial septum-diagonal band of Broca
NBQX	2,3-dioxo-6-nitro-1,2,3,4-tetrahydrobenzo[f]quinoxaline-7-sulfonamide)
NMDA	N-methyl-D-aspartic acid
NMDAR	NMDA receptor
PC	pyramidal cell
PF	parallel fiber
PICK1	protein interacting with C kinase 1
PKC	protein kinase C
PLC	phospholipase C
PN	Purkinje neuron
PSD	postsynaptic density
REM	rapid eye movement
SEM	standard error of mean
Siah-1A	seven in absentia homolog-1A
TBOA	threo-b-benzyloxyaspartic acid
TTX	tetrodotoxin
VFO	very fast oscillation

Curriculum Vitae

Surname	Fan
First name	Wei
Date of birth	27.07.1980
Nationality	China

Education

1998	Graduation from No. 5 high school, Beijing, China
1998 – 2002	College of Life Sciences, Peking University, Beijing, China Bachelor's of Science Thesis title: The activity of <i>crp</i> promoter depends on the coordination of carbon and nitrogen catabolism
2003 – 2006	Kyushu University, Fukuoka, Japan Master's of Systems Life Sciences Thesis title: Isolation and characterization of a CHO <i>pex1</i> mutant with a temperature-sensitive phenotype
2006 – present	Brain Research Institute, University of Zurich, Switzerland employed as PhD student since 06.2006 Thesis title: Conditions for activation of group I metabotropic glutamate receptors in rat hippocampus Thesis committee: Prof. Dr. Stephan Neuhauss Prof. Dr. Urs Gerber Prof. Dr. Peter Sonderegger

Acknowledgements

I would like to express my gratitude to Urs Gerber for inviting me to work in his group, for his continuous support throughout my work and for absolutely great personality.

I would like to thank Stephan Neuhauss and Peter Sonderegger for supporting my thesis as co-supervisors.

I would like to thank Jürg Streit for writing a report on my PhD thesis.

I would like to thank Helge Johannssen for helping me write the summary in German.

I would like to thank Christine Gee for teaching me the theory and the experimental technique of electrophysiology.

I am grateful to Jeanne Ster, Guillaume Coiret, Beat Gähwiler and the members of the Fritjof Helmchen group for valuable discussions.

It would be impossible to accomplish my project without the help of our technical personnel. Lotty Rietschin, Dubravka Göckeritz-Dujmovic, Lubka Spassova prepare hippocampal slice cultures. Hansjörg Kasper and Stefan Giger provide excellent support on electrophysiology device. Roland Schöb and Eva Hochreutener help me make posters.

I thank Neuroscience Center Zurich (ZNZ) for providing me the PhD grant.

I thank all the HIFO members.

Last but not least, I thank all my friends in Switzerland.